

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number
WO 01/60806 A2

(51) International Patent Classification⁷: **C07D 241/00**

(21) International Application Number: **PCT/US01/05264**

(22) International Filing Date: 16 February 2001 (16.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/182,934 16 February 2000 (16.02.2000) US
60/206,455 22 May 2000 (22.05.2000) US

(71) Applicant: **NEUROGEN CORPORATION** [US/US]; 35
Northeast Industrial Road, Branford, CT 06405 (US).

(71) Applicants and

(72) Inventors: **YOON, Taeyoung** [KR/US]; 6 Finch Lane,
Guilford, CT 06437 (US). **GE, Ping** [CN/US]; 718R
Higginum Road, Durham, CT 06422 (US). **HORVATH,**
Raymond, F. [US/US]; 973 Little Meadow, Guilford, CT
06437 (US). **DE LOMBAERT, Stephane** [BE/US]; 37
Concord Drive, Madison, CT 06443 (US). **HODGETTS,**
Kevin, J. [GB/US]; 224 Reservoir Road, Killingworth, CT
06419 (US). **DOLLER, Dario** [US/US]; 20 Killen Road,
Wallingford, CT 06492 (US). **ZHANG, Cunyu** [US/US];
109 Montoya Drive, Branford, CT 06405 (US).

(74) Agent: **CORLESS, Peter, F.**; Edwards & Angell LLP,
Dike, Bronstein, Roberts & Cushman, Intellectual Property
Group, 130 Water Street, Boston, MA 02109 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **SUBSTITUTED ARYLPYRAZINES**

(57) Abstract: Arylpyrazine compounds are provided, including arylpyrazines that can bind with high affinity and high selectivity to CRF₁ receptors, including human CRF₁ receptors. The invention thus includes methods for treatment of disorders and diseases associated with CRF₁ receptors, including CNS-related disorders and diseases, particularly affective disorders and diseases, and acute and chronic neurological disorders and diseases.

WO 01/60806 A2

SUBSTITUTED ARYLPYRAZINES

5 This application claims the benefit of U.S. Provisional Application Serial No.60/182,934 filed February 16, 2000 and of U.S. Provisional Application Serial No.60/206,455 filed May 22, 2000, the teachings of which are incorporated herein by reference.

FIELD OF THE INVENTION

10 The present invention relates to novel arylpyrazine compounds that have useful pharmacological properties in that they bind to cellular receptors, including CRF receptors. Certain of these compounds can bind to and modulate the activity of such receptors. Importantly, the compounds of the invention include compounds that
15 Releasing Factor type 1 Receptors). This invention also relates to pharmaceutical compositions comprising such compounds and to the use of such compounds as pharmaceutical agents, e.g., in treatment of psychiatric disorders and neurological diseases, including major depression, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy and feeding disorders, as well as treatment of
20 immunological, cardiovascular or heart-related diseases and colonic hypersensitivity associated with psychopathological disturbance and stress. Additionally this invention relates to the use such compounds as probes for the localization of cellular receptors in tissue sections.

25 BACKGROUND OF THE INVENTION

 Corticotropin releasing factor (CRF), a 41 amino acid peptide, is the primary physiological regulator of proopiomelanocortin (POMC) derived peptide secretion from the anterior pituitary gland. In addition to its endocrine role at the pituitary gland, immunohistochemical localization of CRF has demonstrated that the hormone
30 has a broad extrahypothalamic distribution in the central nervous system and produces a wide spectrum of autonomic, electrophysiological and behavioral effects consistent with a neurotransmitter or neuromodulator role in brain. There is also evidence that

a significant role in integrating the response of the immune system to physiological, psychological, and immunological stressors.

Clinical data provide evidence that CRF has a role in psychiatric disorders and neurological diseases including depression, anxiety-related disorders and feeding disorders. A role for CRF has also been postulated in the etiology and pathophysiology of Alzheimer's disease, Parkinson's disease, Huntington's disease, progressive supranuclear palsy and amyotrophic lateral sclerosis as they relate to the dysfunction of CRF neurons in the central nervous system.

In affective disorder, or major depression, the concentration of CRF is significantly increased in the cerebral spinal fluid (CSF) of drug-free individuals. Furthermore, the density of CRF receptors is significantly decreased in the frontal cortex of suicide victims, consistent with a hypersecretion of CRF. In addition, there is a blunted adrenocorticotropin (ACTH) response to CRF (i.v. administered) observed in depressed patients. Preclinical studies in rats and non-human primates provide additional support for the hypothesis that hypersecretion of CRF may be involved in the symptoms seen in human depression. There is also preliminary evidence that tricyclic antidepressants can alter CRF levels and thus modulate the numbers of CRF receptors in brain.

CRF has also been implicated in the etiology of anxiety-related disorders. CRF produces anxiogenic effects in animals and interactions between benzodiazepine / non-benzodiazepine anxiolytics and CRF have been demonstrated in a variety of behavioral anxiety models. Preliminary studies using the putative CRF receptor antagonist -helical ovine CRF (9-41) in a variety of behavioral paradigms demonstrate that the antagonist produces "anxiolytic-like" effects that are qualitatively similar to the benzodiazepines. Neurochemical, endocrine and receptor binding studies have all demonstrated interactions between CRF and benzodiazepine anxiolytics providing further evidence for the involvement of CRF in these disorders. Chlordiazepoxide attenuates the "anxiogenic"

effects of CRF in both the conflict test and in the acoustic startle test in rats. The benzodiazepine receptor antagonist Ro 15-1788, which was without behavioral activity alone in the operant conflict test, reversed the effects of CRF in a dose-dependent manner, while the benzodiazepine inverse agonist FG 7142 enhanced the actions of CRF.

5

CRF has also been implicated in the pathogenesis of certain immunological, cardiovascular or heart-related diseases such as hypertension, tachycardia and congestive heart failure, stroke and osteoporosis, as well as in premature birth, psychosocial dwarfism, stress-induced fever, ulcer, diarrhea, post-operative ileus and colonic hypersensitivity associated with psychopathological disturbance and stress.

10

The mechanisms and sites of action through which conventional anxiolytics and antidepressants produce their therapeutic effects remain to be fully elucidated. It has been hypothesized however, that they are involved in the suppression of CRF hypersecretion that is observed in these disorders. Of particular interest are that preliminary studies examining the effects of a CRF receptor antagonist peptide (α -helical CRF₉₋₄₁) in a variety of behavioral paradigms have demonstrated that the CRF antagonist produces "anxiolytic-like" effects qualitatively similar to the benzodiazepines.

15

Certain small molecule compounds for the treatment of CRF related disorders have been disclosed in the literature [for a review see J. McCarthy et al. *Curr. Pharm. Des.* 5: 289 (1999)]. However, none of these compounds has an arylpyrazine structure.

20

Cox et al. (WO 98/38174) have disclosed certain aryl pyrazine derivatives for use as sodium channel blockers in the treatment of central nervous system disorders. The Cox application requires that the arylpyrazine compounds be substituted with two-amino or amido groups.


25

SUMMARY OF THE INVENTION

We have now discovered novel arylpyrazine compounds, including arylpyrazines that can bind with high affinity and high selectivity to CRF₁ receptors, including human CRF₁ receptors. The invention thus includes methods for treatment of disorders and diseases associated with CRF₁ receptors, including CNS-related disorders and diseases, particularly affective disorders and diseases, and acute and chronic neurological disorders and diseases.

Arylpyrazine compounds of the invention are preferably substituted at the 2-ring position by a carbocyclic aryl group such as phenyl, naphthyl and the like, or a heteroaromatic group, particularly a heteroaromatic having a nitrogen ring member such as pyridyl, pyrimidinyl and the like. Arylpyrazine compounds of the invention also are preferably substituted at the 5-ring position (para with respect to the aryl substituent) by a non-hydrogen substituent, particularly a non-aromatic group such as alkyl, alkenyl, alkynyl, heteroalkyl and the like, preferably aminoalkyl and alkoxy. Preferably the 3-ring position of pyrazine compounds of the invention is unsubstituted (i.e. hydrogen) or substituted by other than amino (-NH₂) or alkylamide (-NHC(=O)alkyl, particularly -NHC(=O)C₁₋₄alkyl or -NHC(=O)C₃₋₇cycloalkyl), and/or the 5-ring position of pyrazine compounds of the invention are other than hydrogen, alkyl, aminoalkyl or a nitrogen-containing heteroalicyclic compound.

Typically preferred compounds of the invention include those of the following Formula I:



I

wherein:


Ar is substituted phenyl, optionally substituted naphthyl, or an optionally substituted heterocyclic group having from 1 to 3 rings, and 3 to 8 ring members in each ring and 1 to about 3 hetero atoms;

5 R_1 and R_3 are each independently hydrogen, halogen, cyano, nitro, amino, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted mono or dialkylamino, optionally substituted alkoxy, optionally substituted alkylthio, optionally substituted alkylsulfinyl, or optionally substituted alkylsulfonyl; and

10 R_2 is halogen, cyano, nitro, amino, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkylamino, optionally substituted alkoxy, optionally substituted alkylthio, optionally substituted alkylsulfinyl, or optionally substituted alkylsulfonyl;

15 with the proviso that if Ar is phenyl substituted with halogen, naphthyl, or naphthyl substituted with halogen, then the compounds where R_3 is hydrogen or amino are excluded.

Preferred compounds of the invention also include those of the following Formula IA:



Formula IA

or a pharmaceutically acceptable salt thereof, wherein:

25 R_1 is selected from H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, halogen, CN, C_{1-4} haloalkyl, trifluoromethyl, trifluoromethoxy, $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), $-O(C_{1-4}$ alkyl), and $S(O)_n(C_{1-4}$ alkyl);

R_2 is selected from the group consisting of $-XR_A$ and Y, wherein $-X$, R_A , and Y are defined below; and

R_3 is selected from the group consisting of hydrogen, halogen, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), and $-S(O)_n(C_{1-4}$ alkyl), haloalkyl, trifluoromethyl, trifluoromethoxy, $-XR_A$ and Y;

R_4 is absent or an oxygen atom;

5 Ar is phenyl, mono-, di-, or tri-substituted with R_C , or

Ar is selected from the group consisting of:

naphthyl, pyridyl, pyridonyl, pyrimidinyl, and thiophenyl, each of which is unsubstituted or mono-, di-, or tri-substituted with R_C ;

with the proviso that if Ar is phenyl substituted with halogen, naphthyl, or naphthyl
10 substituted with halogen, then the compounds where R_3 is hydrogen are excluded;

R_A and R_B , which may be the same or different, are independently selected at each occurrence from the group consisting of: hydrogen and straight, branched, or cyclic alkyl groups consisting of 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or
15 more substituent(s) selected from oxo, hydroxy, halogen, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), $-NHC(O)(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl) $C(=O)(C_{1-4}$ alkyl), $-NHS(O)_n(C_{1-4}$ alkyl), $-S(O)_n(C_{1-4}$ alkyl), $-S(O)_nNH(C_{1-4}$ alkyl), $-S(O)_nN(C_{1-4}$ alkyl)(C_{1-4} alkyl), and Z;

R_C is independently selected at each occurrence from the group consisting of halogen,
20 cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C_{1-6} alkyl optionally substituted with 0-2 R_D , C_{2-6} alkenyl substituted with 0-2 R_D , C_{1-4} alkynyl substituted with 0-2 R_D , C_{3-7} cycloalkyl substituted with 0-2 R_D , (C_{3-7} cycloalkyl) C_{1-4} alkyl substituted with 0-2 R_D , $-O(C_{1-4}$ alkyl) substituted with 0-2 R_D , $-NH(C_{1-4}$ alkyl) substituted with 0-2 R_D , $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl) each
25 independently substituted with 0-2 R_D , $-XR_A$, and Y;


R_D is independently selected at each occurrence the group consisting of halogen, hydroxy, cyano, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), morpholino, pyrrolidino, piperidino, thiomorpholino, piperazino, 4-hydroxypiperidino, $-S(O)_n(C_{1-4}$ alkyl), trifluoromethyl, trifluoromethoxy, $CO(C_{1-4}$ alkyl), $CONH(C_{1-4}$ alkyl), $CON(C_{1-4}$ alkyl)(C_{1-4} alkyl), $-XR_A$, and Y;
30

X is independently selected at each occurrence from the group consisting of $-\text{CH}_2-$, $-\text{CHR}_B-$, $-\text{O}-$, $-\text{C}(=\text{O})-$, $-\text{C}(=\text{O})\text{O}-$, $-\text{S}(\text{O})_n-$, $-\text{NH}-$, $-\text{NR}_B-$, $-\text{C}(=\text{O})\text{NH}-$, $-\text{C}(=\text{O})\text{NR}_B-$, $-\text{S}(\text{O})_n\text{NH}-$, $-\text{S}(\text{O})_n\text{NR}_B-$, $-\text{OC}(=\text{S})\text{S}-$, $-\text{NHC}(=\text{O})-$, $-\text{NR}_B\text{C}(=\text{O})-$, $-\text{NHS}(\text{O})_n-$, $-\text{OSiH}_n(\text{C}_{1-4}\text{alkyl})_{2-n}-$, and $-\text{NR}_B\text{S}(\text{O})_n-$; and

- 5 Y and Z are independently selected at each occurrence from the group consisting of: 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents selected from halogen, haloalkyl, oxo, hydroxy, amino, C_{1-4} alkyl, $-\text{O}(\text{C}_{1-4}$ alkyl), $-\text{NH}(\text{C}_{1-4}$ alkyl), $-\text{N}(\text{C}_{1-4}$ alkyl)(C_{1-4} alkyl), and $-\text{S}(\text{O})_n(\text{C}_{1-4}$ alkyl), and
- 10 said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) selected from N, O, and S, with the point of attachment being either carbon or nitrogen; and

n is independently selected at each occurrence from 0, 1, and 2.

- 15 Particularly preferred arylpyrazines of the invention include compounds of the following Formula IB:



wherein:

- 20 Ar is selected from the group consisting of phenyl, 1- or 2-naphthyl, 2-, 3-, or 4-pyridyl, 2-, 4-, or 5-pyrimidinyl, each Ar optionally substituted with 1 to 5 with R_6 group, with the proviso that at least one of the positions ortho or para to the point of attachment of Ar to the pyrazine ring system is substituted;
- R₁ is selected from H, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, $(\text{C}_{3-6}$ cycloalkyl) C_{1-4} alkyl,
- 25 halogen, CN, C_{1-4} haloalkyl, $-\text{NH}(\text{C}_{1-4}$ alkyl), $-\text{N}(\text{C}_{1-4}$ alkyl)(C_{1-4} alkyl), $-\text{O}(\text{C}_{1-4}$ alkyl), and $-\text{S}(\text{O})_b(\text{C}_{1-4}$ alkyl), wherein b is 0, 1, or 2;
- R₂ is selected from the group consisting of $-\text{XR}_4$ and Y, wherein X, R₄, and Y are as defined below;

X is independently selected at each occurrence from the group consisting of $-\text{CH}_2-$, $-\text{CHR}_5-$, $-\text{O}-$, $-\text{S}(\text{O})_n-$, $-\text{NH}-$, $-\text{NR}_5-$, $-\text{C}(=\text{O})\text{NH}-$, $-\text{C}(=\text{O})\text{NR}_5-$, $-\text{S}(\text{O})_n\text{NH}-$, $-\text{S}(\text{O})_n\text{NR}_5-$, $-\text{NHC}(=\text{O})-$, $-\text{NR}_5\text{C}(=\text{O})-$, $-\text{NHS}(\text{O})_n-$, and $-\text{NR}_5\text{S}(\text{O})_n-$ (where n is 0, 1, or 2);

- 5 Y and Z are independently selected at each occurrence from the group consisting of:
 3- to 7-membered heterocycles, saturated or unsaturated, containing one or more heteroatom(s) selected from N, O, and S, with the point of attachment being either carbon or nitrogen (where applicable), and which may be further substituted with one or more substituents selected from halogen, oxo, hydroxy, amino, C_{1-4} alkyl, $-\text{O}(\text{C}_{1-4}$ alkyl),
 10 $-\text{NH}(\text{C}_{1-4}$ alkyl), $-\text{N}(\text{C}_{1-4}$ alkyl)(C_{1-4} alkyl), and $-\text{S}(\text{O})_a(\text{C}_{1-4}$ alkyl) (wherein a is 0, 1, or 2);

R_3 is selected from the group consisting of hydrogen, halogen, C_{1-4} alkyl, $-\text{O}(\text{C}_{1-4}$ alkyl), $-\text{NH}(\text{C}_{1-4}$ alkyl), $-\text{N}(\text{C}_{1-4}$ alkyl)(C_{1-4} alkyl), and $-\text{S}(\text{O})_c(\text{C}_{1-4}$ alkyl) (wherein c is 0, 1, or 2), trifluoromethyl, trifluoromethoxy, $-\text{XR}_4$, and Y, where
 15 $-\text{X}$, R_4 , and Y are as defined above.

R_4 and R_5 , which may be the same or different, are independently selected at each occurrence from the group consisting of:
 hydrogen, straight, branched, or cyclic alkyl groups consisting of 1 to 8 carbon atoms,
 20 which may contain one or more double or triple bonds, each of which may optionally be further substituted with one or more substituent(s) selected from oxo, hydroxy, $-\text{O}(\text{C}_{1-4}$ alkyl), $-\text{NH}(\text{C}_{1-4}$ alkyl), $-\text{N}(\text{C}_{1-4}$ alkyl)(C_{1-4} alkyl), $-\text{NHC}(\text{O})(\text{C}_{1-4}$ alkyl), $-\text{N}(\text{C}_{1-4}$ alkyl) $\text{C}(\text{O})(\text{C}_{1-4}$ alkyl), $-\text{NHS}(\text{O})_m(\text{C}_{1-4}$ alkyl), $-\text{S}(\text{O})_m(\text{C}_{1-4}$ alkyl), $-\text{S}(\text{O})_m\text{NH}(\text{C}_{1-4}$ alkyl), $-\text{S}(\text{O})_m\text{N}(\text{C}_{1-4}$ alkyl)(C_{1-4} alkyl), (where m is 0, 1, or 2), and Z, wherein Z is as
 25 defined above;

R_6 is independently selected at each occurrence from halogen, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C_1 - C_6 alkyl (optionally substituted with 0-2 R_7 , C_{2-6} alkenyl substituted with 0-2 R_7 , C_{1-4} alkynyl substituted with 0-2 R_7 , C_{3-6} cycloalkyl substituted with 0-2 R_7 , $(\text{C}_{3-6}$ cycloalkyl) C_{1-4} alkyl substituted with 0-2 R_7 ,
 30 $\text{O}(\text{C}_{1-4}$ alkyl) substituted with 0-2 R_7 , $-\text{NH}(\text{C}_{1-4}$ alkyl) substituted with 0-2 R_7 , $-\text{N}(\text{C}_{1-4}$

alkyl)(C₁₋₄ alkyl) each independently substituted with 0-2 R₇, -XR₄, cyano, and Y;
and

R₇ is independently selected at each occurrence the group consisting of halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), morpholino,
5 pyrrolidino, piperidino, thiomorpholino, piperazino, 4-hydroxypiperidino, -S(O)_p(C₁₋₄ alkyl), trifluoromethyl, trifluoromethoxy, CO(C₁₋₄alkyl), CONH(C₁₋₄ alkyl), CON(C₁₋₄ alkyl)(C₁₋₄ alkyl), -XR₄, and Y, wherein X, R₄ and Y are defined as above and p is 0, 1 or 2;

and pharmaceutically acceptable salts thereof;

10 and preferably if Ar is phenyl substituted with halogen, naphthyl, or naphthyl substituted with halogen, then the exclusion of compounds where 1) R₃ being hydrogen, amino (-NH₂) or alkylamide particularly -NHC(=O)C₁₋₄alkyl or -NHC(=O)C₃₋₇cycloalkyl, and/or 2) R₂ being hydrogen, alkylamide particularly -NHC(=O)C₁₋₄alkyl or -NHC(=O)C₃₋₇cycloalkyl, amino substituted by one or two
15 alkyl particularly C₁₋₄ alkyl, or a nitrogen-containing 6-membered ring.

In the above Formulae I, IA, and IB, as well as below-specified Formulae Ia through Ie,

II, IIa through Iii, III, IIIa through IIIe and IV, preferred Ar groups include 2,4-dichlorophenyl, 2,4-dimethoxyphenyl, 4-methoxy-2-methylphenyl, 2,4,6-trimethylphenyl, 4-methoxy-2,6-dimethylphenyl, 4-cyano-2,6-dimethylphenyl, 2-methoxy-4,6-dimethylphenyl, 2-methoxy-4,6-bis(trifluoromethyl)phenyl, 2,6-dichloro-4-methoxyphenyl, 2-(2-(1-morpholino)ethoxy)-4,6-dimethylphenyl, 2-(2-(4-hydroxy-1-piperidino)ethoxy)-4,6-dimethylphenyl, 2-methyl-4-(dimethylamino)-3-pyridyl, and 2-chloro-4-dimethylamino-3-pyridyl, 2-methoxy-
20 4,6-dimethyl-3-pyridyl.

In the above Formulae I, IA, and IB, as well as below-specified Formulae Ia through Ie, II, IIa through Iii, III, IIIa through IIIe and IV, preferred R₁ groups include methyl, ethyl, chloro, bromo, iodo, trifluoromethyl, methoxy, dimethylamino, and
30 thiomethoxy.

In the above Formulae I, IA, and IB, as well as below-specified Formulae Ia through Ie, II, IIa through III, IIIa through IIIe and IV, preferred R₂ groups include dipropylamino, bis(2-methoxyethyl)amino, 4-methyl-1-piperazino, 3-pentylamino, 4-heptylamino, 1,3-dimethoxy-2-propylamino, 1-(dimethylamino)-2-pentylamino, 1-(3-pyridyl)-2-butylamino, (2-methoxy-5-pyridine)amino, 3-pentyloxy, 4-heptyloxy, 1-methoxy-2-butyloxy, and 1,3-dimethoxy-2-propyloxy.

In the above Formulae I, IA, and IB, as well as below-specified Formulae Ia through Ie, II, IIa through III, IIIa through IIIe and IV, preferred R₃ groups include methyl, ethyl, chloro, trifluoromethyl, methoxy, dimethylamino, thiomethoxy, methanesulfonyl, (1-morpholino)methyl, 2-(1-pyrrolidino)ethyl, and (2-methoxy)ethoxy.

Preferred arylpyrazines of the invention exhibit good activity in a standard in vitro CRF receptor binding assays, specifically the assay as specified in Example 96 which follows. Particularly preferred arylpyrazines of the invention have an IC₅₀ in such a defined standard in vitro CRF receptor binding assay of about 1.5 micromolar or less, still more preferably an IC₅₀ of about 100 nanomolar or less, or even more preferably an IC₅₀ of about 10 nanomolar or less or even 1 nanomolar or less in a such a defined standard in vitro CRF receptor binding assay.

Preferred arylpyrazines of the invention do not show activity in a standard in vitro Na channel functional assay, specifically the assay as specified in Example 99 which follows. Particularly preferred arylpyrazines of the invention do not show any statistically significant activity in a defined standard in vitro Na channel functional assay 0 at the p < 0.05 level of significance.

The invention further comprises methods of treating patients suffering from or susceptible to (i.e. prophylactic treatment) certain disorders or diseases with an effective amount of a compound of the invention. The patient may be a human or other mammal

such as other primates. Treatment of humans, domesticated companion animals (pet) or livestock animals suffering from certain with an effective amount of a compound of the invention is encompassed by the invention.

5 Additionally this invention relates to the use of the CRF1 compounds of the invention as probes for the localization of receptors in tissue sections.

 The compounds of the present invention are useful for the treatment of CNS disorders particularly affective disorders, anxiety disorders, stress-related disorders,
10 eating disorders and substance abuse. Affective disorders include all types of depression, bipolar disorder, cyclothymia, and dysthymia. Anxiety disorders include generalized anxiety disorder, panic, phobias and obsessive-compulsive disorder. Stress-related disorders include post-traumatic stress disorder, hemorrhagic stress, stress-induced psychotic episodes, psychosocial dwarfism, stress headaches, stress-induced immune
15 systems disorders such as stress-induced fever, and stress-related sleep disorders. Eating disorders include anorexia nervosa, bulimia nervosa, and obesity. These conditions are herein after referred to as the primary CRF-related CNS disorders.

 Arylpyrazine compounds of the invention (which includes compounds of the
20 Formulae I, IA, and IB, as well as below-specified Formulae Ia through Ie, II, IIa through Iii, III, IIIa through IIIe and IV,) are also useful in the treatment of a variety of neurological disorders including supranuclear palsy, AIDS related dementias, multiinfarct dementia, neurodegenerative disorders such as Alzheimer's disease, Parkinson's disease, and Huntington's disease, head trauma, spinal cord trauma, ischemic neuronal damage,
25 amyotrophic lateral sclerosis, disorders of pain perception such as fibromyalgia and epilepsy. These conditions are hereinafter referred to as the secondary CRF-related CNS disorders.

 Arylpyrazine compounds of the invention are useful as modulators of the G-
30 protein coupled receptor function.

Additionally, arylpyrazine compounds of the invention are useful as modulators of the CRF receptor in the treatment of a number of gastrointestinal, cardiovascular, hormonal, autoimmune and inflammatory conditions. Such conditions include irritable
5 bowel syndrome, ulcers, Crohn's disease, spastic colon, diarrhea, post operative ilius and colonic hypersensitivity associated with psychopathological disturbances or stress, hypertension, tachycardia, congestive heart failure, infertility, euthyroid sick syndrome, inflammatory conditions effected by rheumatoid arthritis and osteoarthritis, pain, asthma, psoriasis and allergies. These conditions are referred to hereinafter as the non-CNS CRF-
10 related disorders.

Arylpyrazine compounds of the invention are also useful as modulators of the CRF₁ receptor in the treatment of animal conditions associated with aberrant CRF levels. These conditions include porcine stress syndrome, bovine shipping fever, equine
15 paroxysmal fibrillation, and dysfunctions induced by confinement in chickens, sheering stress in sheep or human-animal interaction related stress in dogs, psychosocial dwarfism and hypoglycemia. These animal conditions are referred to hereinafter as the CRF-related animal disorders.

20 According to yet another aspect, the present invention provides pharmaceutical compositions comprising one or more aryl pyrazine compound of the invention of a pharmaceutically acceptable salt thereof, particularly for use in the treatment of any of the primary CNS disorders, secondary CNS disorders, non-CNS CRF-related disorders, or CRF-related animal disorders.


25

According to a still further aspect of the invention, arylpyrazine compounds of the invention (and especially labeled compounds of this invention) are useful as standards and reagents in determining the ability of a potential pharmaceutical to bind to the CRF
30 receptor.

Other aspects of the invention are disclosed *infra*.

DETAILED DESCRIPTION OF THE INVENTION

The invention provides compounds of Formula I shown above. Preferred
 5 compound of the invention include those of the following Formula Ia:



or a pharmaceutically acceptable salt thereof, wherein:

R₁ is selected from H, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, halogen, CN, C₁₋₄ haloalkyl, trifluoromethyl, trifluoromethoxy, -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), -
 10 O(C₁₋₄ alkyl), and S(O)_n(C₁₋₄ alkyl);

R₂ is selected from the group consisting of -XR_A and Y, wherein -X, R_A, and Y are defined below; and

R₃ is selected from the group consisting of hydrogen, halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), and -S(O)_n(C₁₋₄ alkyl), haloalkyl,
 15 trifluoromethyl, trifluoromethoxy, -XR_A and Y;

R₄ is absent or an oxygen atom;

Ar is phenyl, mono-, di-, or tri-substituted with R_C, or

Ar is selected from the group consisting of:
 naphthyl, pyridyl, pyridonyl, pyrimidinyl, and thiophenyl, each of which is
 20 unsubstituted or mono-, di-, or tri-substituted with R_C;

with the proviso that if Ar is phenyl substituted with halogen, naphthyl, or naphthyl substituted with halogen, then the compounds where R₃ is hydrogen are excluded;

R_A and R_B, which may be the same or different, are independently selected at each occurrence from the group consisting of: hydrogen and straight, branched, or
 25 cyclic alkyl groups consisting of 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) selected from oxo, hydroxy, halogen, -O(C₁₋₄ alkyl), -NH(C₁₋₄

alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), -NHC(O)(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)C(=O)(C₁₋₄ alkyl), -NHS(O)_n(C₁₋₄ alkyl), -S(O)_n(C₁₋₄ alkyl), -S(O)_nNH(C₁₋₄ alkyl), -S(O)_nN(C₁₋₄ alkyl)(C₁₋₄ alkyl), and Z;

5 R_C is independently selected at each occurrence from the group consisting of halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl optionally substituted with 0-2 R_D, C₂₋₆ alkenyl substituted with 0-2 R_D, C₁₋₄ alkynyl substituted with 0-2 R_D, C₃₋₇ cycloalkyl substituted with 0-2 R_D, (C₃₋₇ cycloalkyl)C₁₋₄ alkyl substituted with 0-2 R_D, -O(C₁₋₄ alkyl) substituted with 0-2 R_D, -NH(C₁₋₄ alkyl) substituted with 0-2 R_D, -N(C₁₋₄ alkyl)(C₁₋₄ alkyl) each
10 independently substituted with 0-2 R_D, -XR_A, and Y;

R_D is independently selected at each occurrence the group consisting of halogen, hydroxy, cyano, C₁₋₄alkyl, -O(C₁₋₄alkyl), -NH(C₁₋₄alkyl), -N(C₁₋₄alkyl)(C₁₋₄alkyl), morpholino, pyrrolidino, piperidino, thiomorpholino, piperazino, 4-hydroxypiperidino, -S(O)_n(C₁₋₄alkyl), trifluoromethyl, trifluoromethoxy, CO(C₁₋₄alkyl), CONH(C₁₋₄alkyl), CON(C₁₋₄alkyl)(C₁₋₄alkyl), -XR_A, and Y;
15

X is independently selected at each occurrence from the group consisting of -CH₂-, -CHR_B-, -O-, -C(=O)-, -C(=O)O-, -S(O)_n-, -NH-, -NR_B-, -C(=O)NH-, -C(=O)NR_B-, -S(O)_nNH-, -S(O)_nNR_B-, -OC(=S)S-, -NHC(=O)-, -NR_BC(=O)-, -NHS(O)_n-, -OSiH_n(C₁₋₄alkyl)_{2-n}-, and -NR_BS(O)_n-; and

20 Y and Z are independently selected at each occurrence from the group consisting of: 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents selected from halogen, haloalkyl, oxo, hydroxy, amino, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), and -S(O)_n(C₁₋₄ alkyl), and
25 said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) selected from N, O, and S, with the point of attachment being either carbon or nitrogen; and

n is independently selected at each occurrence from 0, 1, and 2.

30 Preferred compounds of general Formula Ia are those where:

R_4 is absent and Ar is phenyl, mono-, di-, or tri-substituted with R_C , and R_1 and R_3 are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy and methoxy.

Such compounds are referred to herein as compounds of general Formula Ib.

5

More preferred compounds of the invention are those of general Formula Ia where:

R_4 is absent and Ar is phenyl, mono-, di-, or tri-substituted with R_C ; and R_A and R_B , which may be the same or different, are independently straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds.

10

Such compounds are referred to as compounds of Formula Ic.

15

Particularly preferred are compounds of general Formula Ia are those where:

R_4 is absent and Ar is phenyl mono-, di-, or tri-substituted with R_C ; R_A and R_B , which may be the same or different, are independently selected at each occurrence from the group consisting of: straight, branched, and cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds; and


R_1 and R_3 are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy, and methoxy.

20

Such compounds are referred to herein as compounds of general Formula Id.

Particularly preferred are compounds of general Formula Ia are those where:


25



wherein A is NR_A or O.

Such compounds are referred to herein as compounds of general Formula Ie.

Even more preferred compounds of the invention are those of general Formula II:



5 Formula II

wherein

R_X and R_Y are the same or different and are independently selected straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more
 10 double or triple bonds, each of which may be further substituted with one or more substituent(s) independently selected from hydroxy, halogen, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), and optionally substituted phenyl.

Other preferred compounds of the invention include those of general Formula II (above)
 15 and pharmaceutically acceptable salts thereof, wherein:

Ar is phenyl, mono-, di-, or tri-substituted with R_C; and R₁ and R₃ are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy and methoxy.

Such compounds are referred to herein as compounds of general Formula IIa.

20

Other preferred compounds of the invention include those of general Formula II (above) and pharmaceutically acceptable salts thereof, wherein:

R₁ and R₃ are independently chosen from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and
 25 trifluoromethoxy; and

Ar is phenyl, which is mono-, di-, or trisubstituted with one or more substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.

Such compounds are referred to herein as compounds of general Formula IIb.

5

Other preferred compounds of the invention include those of general Formula II (above) and pharmaceutically acceptable salts thereof, wherein:

R_x is hydrogen;

R_y is chosen from the group consisting of:

10 straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds;

R₁ and R₃ are independently chosen from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

15 Ar is phenyl, which is mono-, di-, or trisubstituted with substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.

20 Such compounds are referred to herein as compounds of general Formula IIc.

Other preferred compounds of the invention include those of general Formula II (above) and pharmaceutically acceptable salts thereof, wherein:


R_x is hydrogen;

25 R_y is chosen from the group consisting of:

straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds;

R₁ and R₃ are independently chosen from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and
30 trifluoromethoxy; and

Ar is a phenyl group of the formula:



wherein L indicates a bond to the pyrazine ring in Formula A;

and the phenyl group is substituted at one, two or three of positions 2, 4, and 6 with
 5 substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino,
 and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄
 alkoxy), and mono- or di(C₁₋₄ alkyl)amino.

Such compounds are referred to herein as compounds of general Formula IId.

10

Other preferred compounds of the invention include those of general Formula II (above)
 and pharmaceutically acceptable salts thereof, wherein:


R_x is hydrogen;

R_y is chosen from the group consisting of:

15 straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may
 contain one or more double or triple bonds;

R₁ and R₃ are independently chosen from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄
 alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and
 trifluoromethoxy; and

20 Ar is a phenyl group of the formula:



wherein L indicates a bond to the pyrazine ring in Formula A;


and the phenyl group is substituted at positions 2 and 4 with substituents independently
 selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.

Such compounds are referred to herein as compounds of general Formula IIe.

5


Other preferred compounds of the invention include those of general Formula II (above) and pharmaceutically acceptable salts thereof, wherein:



wherein:

- 10 R₁ and R₃ are independently chosen at each occurrence from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and


Ar is a phenyl group of the formula:



- 15 wherein L indicates a bond to the pyrazine ring in Formula A;
and the phenyl group is substituted at one, two or three of positions 2, 4, and 6 with substituent(s) independently selected from:
halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.
- 20

Such compounds are referred to herein as compounds of general Formula IIf.

Other preferred compounds of the invention include those of general Formula II (above) and pharmaceutically acceptable salts thereof, wherein:



wherein

R is independently selected at each occurrence from the group consisting of: hydrogen, halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, C₁₋₆alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino; and

R₁ and R₃ are independently chosen from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy.

Such compounds are referred to herein as compounds of general Formula IIg.

Other preferred compounds of the invention include those of general Formula II (above) and pharmaceutically acceptable salts thereof, wherein:


Ar is phenyl, which is mono-, di-, or tri-substituted with R_C;

R_X and R_Y, which may be the same or different, are independently selected at each occurrence from the group consisting of: straight, branched, and cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds; and


R₁ and R₃ are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy, and methoxy.

Such compounds are referred to herein as compounds of general Formula IIh.

Other preferred compounds of the invention include those of general Formula II (above) and pharmaceutically acceptable salts thereof, wherein:




wherein R_x and R_y are independently hydrogen or C_{1-8} alkyl; or NR_xR_y is:



wherein z is 0 or 1.

Such compounds are referred to herein as compounds of general Formula Iii.

10 Other highly preferred compounds are those of general Formula III:



III

wherein:

R_x is chosen from the group consisting of:

- straight, branched, or cyclic alkyl groups, including (cycloalkyl)alkyl groups, having
- 15 from 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) independently selected from
- (a) hydroxy, halogen, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), and
 - 20 (b) 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents selected from halogen, haloalkyl, oxo, hydroxy, amino,

C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), and $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl) and wherein said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) selected from N, O, and S, with the point of attachment being either carbon or nitrogen.

5

Other preferred compounds of the invention include those of general Formula III (above) and pharmaceutically acceptable salts thereof, wherein:

R_x is selected from straight, branched, or cyclic alkyl groups containing of 1 to 8 carbon atoms, which may contain one or more double or triple bonds;

10 and;

R_1 and R_2 are independently chosen from halogen, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

Ar is phenyl, which is mono-, di-, or trisubstituted with one or more substituent(s) independently selected from:


halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy(C_{1-4} alkoxy), mono- or di(C_{1-4})amino(C_{1-4} alkoxy), and mono- or di(C_{1-4} alkyl)amino.

Such compounds are referred to herein as compounds of general Formula IIIa.

20

Other preferred compounds of the invention include those of general Formula III (above) and pharmaceutically acceptable salts thereof, wherein:

Ar is a phenyl group of the formula:



25 wherein L indicates a bond to the pyrazine ring in Formula III;

and the phenyl group is substituted at one, two, or three of positions 2, 4, and 6 with substituent(s) independently selected from:


halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.

Such compounds are referred to herein as compounds of general Formula IIIb.

5

Other preferred compounds of the invention include those of general Formula III (above) and pharmaceutically acceptable salts thereof, wherein:

Ar is a phenyl group of the formula:




10 wherein L indicates a bond to the pyrazine ring in Formula III;

and the phenyl group is substituted at positions 2 and 4 with substituents independently selected from halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.

15 Such compounds are referred to herein as compounds of general Formula IIIc.


Other preferred compounds of the invention include those of general Formula III (above) and pharmaceutically acceptable salts thereof, wherein:



20

R₁ and R₃ are independently chosen from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

Ar is a phenyl group of the formula:



wherein L indicates a bond to the pyrazine ring in Formula B;

and the phenyl group is substituted at one, two, or three of positions 2, 4, and 6 with
 5 substituent(s) independently selected from: halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄alkyl)amino.

Such compounds are referred to herein as compounds of general Formula IIIId.


10 Other preferred compounds of the invention include those of general Formula III (above) and pharmaceutically acceptable salts thereof, wherein:

R₁ and R₃ are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy and methoxy.

Such compounds are referred to herein as compounds of general Formula IIIe.

15

Another preferred compound of the invention include 2-(2-methoxy-5-trifluoromethoxyphenyl)-3-ethyl-6-methylamino-5-(1-ethylpropoxy)-pyrazine.
 and pharmaceutically acceptable salts thereof, the compound structure is:



20

DEFINITIONS

The compounds herein described may have one or more asymmetric centers or planes. Compounds of the present invention containing an asymmetrically substituted

atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms (racemates), by asymmetric synthesis, or by synthesis from optically active starting materials. Resolution of the racemates can be accomplished, for example, by conventional methods such as

5 crystallization in the presence of a resolving agent, or chromatography, using, for example a chiral HPLC column. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. *Cis* and *trans* geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of

10 isomers or as separated isomeric forms. All chiral (enantiomeric and diastereomeric), and racemic forms, as well as all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

The CRF antagonist compounds provided by this invention and labeled

15 derivatives thereof are also useful as standards and reagents in determining the ability of a potential pharmaceutical to bind to the CRF receptor.

Labeled derivatives the CRF antagonist compounds provided by this invention are also useful as radiotracers for positron emission tomography (PET) imaging or for single

20 photon emission computerized tomography (SPECT).

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valence is not exceeded, and that the substitution results in a

25 stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties. The present invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers, By way of general example and without limitation, isotopes of hydrogen include tritium and


30 deuterium. Isotopes of carbon include ^{11}C , ^{13}C , and ^{14}C .

When any variable occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R*, then
5 said group may optionally be substituted with up to two R* groups and R* at each occurrence is selected independently from the definition of R*. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

10 As indicated above, various substituents of the various formulae are "optionally substituted", including Ar, R₁, R₂, and R₃ of Formula I, and such substituents as recited in the sub-formulae such as Formulae Ia and the like. When substituted, those substituents (Ar, R₁, R₂, and R₃) may be substituted by other than hydrogen at one or more available positions, typically 1 to 3 or 4 positions, by one or more suitable groups such as those
15 disclosed herein. Suitable groups that may be present on a "substituted" Ar, R₁, R₂, and R₃ group or other substituent include e.g. halogen such as fluoro, chloro, bromo and iodo; cyano; hydroxyl; nitro; azido; alkanoyl such as a C₁₋₆ alkanoyl group such as acyl and the like; carboxamido; alkyl groups including those groups having 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5, or 6 carbon atoms; alkenyl and alkynyl groups including groups
20 having one or more unsaturated linkages and from 2 to about 12 carbon, or 2, 3, 4, 5 or 6 carbon atoms; alkoxy groups having those having one or more oxygen linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5 or 6 carbon atoms; aryloxy such as phenoxy; alkylthio groups including those moieties having one or more thioether linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5 or 6 carbon atoms; alkylsulfinyl groups
25 including those moieties having one or more sulfinyl linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5, or 6 carbon atoms; alkylsulfonyl groups including those moieties having one or more sulfonyl linkages and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5, or 6 carbon atoms; aminoalkyl groups such as groups having one or more N atoms and from 1 to about 12 carbon atoms, or 1, 2, 3, 4, 5 or 6 carbon atoms; carbocyclic
30 aryl having 6 or more carbons, particularly phenyl (e.g. an Ar group being a substituted or

unsubstituted biphenyl moiety); aralkyl having 1 to 3 separate or fused rings and from 6 to about 18 carbon ring atoms, with benzyl being a preferred group; aralkoxy having 1 to 3 separate or fused rings and from 6 to about 18 carbon ring atoms, with O-benzyl being a preferred group; or a heteroaromatic or heteroalicyclic group having 1 to 3 separate or fused rings with 3 to about 8 members per ring and one or more N, O or S atoms, e.g. coumarinyl, quinolinyl, pyridyl, pyrazinyl, pyrimidyl, furyl, pyrrolyl, thienyl, thiazolyl, oxazolyl, imidazolyl, indolyl, benzofuranyl, benzothiazolyl, tetrahydrofuranyl, tetrahydropyranyl, piperidinyl, morpholino and pyrrolidinyl.

As used herein, the substitution position of a substituent on a phenyl ring is dependent on the point of attachment of the substituent to the phenyl ring relative to the point of attachment of the phenyl group to the remainder of the chemical compound. For example, L indicates the point of attachment of the phenyl ring to the remainder of the chemical compound. The numbers 2, 3, 4, 5 and 6 identify the individual ring atoms to which substituents can be attached.



As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups, having the specified number of carbon atoms. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, and s-pentyl. Alkyl groups typically have 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms. Preferred alkyl groups are C₁-C₆ alkyl groups. Especially preferred alkyl groups are methyl, ethyl, propyl, butyl, 3-pentyl.

As used herein, "cycloalkyl" is intended to include saturated ring groups, having the specified number of carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl. Cycloalkyl groups typically will have 3 to about 8 ring members.

5 In the term "(C₃₋₆ cycloalkyl)C₁₋₄ alkyl", as defined above, the point of attachment is on the alkyl group. This term encompasses, but is not limited to, cyclopropylmethyl, cyclohexylmethyl, cyclohexylmethyl.

10 As used here, "alkenyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl and propenyl. Alkenyl groups typically will have 2 to about 12 carbon atoms, more typically 2 to about 12 carbon atoms.

15 As used herein, "alkynyl" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more triple carbon-carbon bonds which may occur in any stable point along the chain, such as ethynyl and propynyl. Alkynyl groups typically will have 2 to about 16 carbon atoms, more typically 2 to about 12 carbon atoms.

20 As used herein, "haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example -C_vF_w where v = 1 to 3 and w = 1 to (2v+1)). Examples of haloalkyl include, but are not limited to, 25 trifluoromethyl, trichloromethyl, pentafluoroethyl, and pentachloroethyl. Typical haloalkyl groups will have 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms.

30 As used herein, "alkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of

alkoxy include, but are not limited to, methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, 2-butoxy, t-butoxy, n-pentoxy, 2-pentoxy, 3-pentoxy, isopentoxy, neopentoxy, n-hexoxy, 2-hexoxy, 3-hexoxy, and 3-methylpentoxy. Alkoxy groups typically have 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms.

5

As used herein, the term "alkylthio" includes those groups having one or more thioether linkages and suitably from 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms, still more typically 1 to about 6 or 8 carbon atoms.

10

As used herein, the term "alkylsulfinyl" includes those groups having one or more sulfoxide (SO) linkage groups and suitably from 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms, still more typically 1 to about 6 or 8 carbon atoms.

15

As used herein, the term "alkylsulfonyl" includes those groups having one or more sulfonyl (SO₂) linkage groups and suitably from 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms, still more typically 1 to about 6 or 8 carbon atoms.

20

As used herein, the term "alkylamino" includes those groups having one or more primary, secondary and/or tertiary amine groups and suitably from 1 to about 16 carbon atoms, more typically 1 to about 12 carbon atoms, still more typically 1 to about 6 or 8 carbon atoms.

25

As used herein, "halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo; and "counter-ion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, and the like.

30

As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3- to 7-membered monocyclic or bicyclic or 7- to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl,

cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane, [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, and tetrahydronaphthyl.

5 As used herein, the term "heterocycle" or "heterocycloalkyl" is intended to mean a stable 5-to 7-membered monocyclic or bicyclic or 7-to 10-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the
10 above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be
15 quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" or similar term such as "heteroaryl" is intended to mean a stable 5-to 7-membered monocyclic or bicyclic or 7-to 10-membered bicyclic
20 heterocyclic aromatic ring which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the aromatic heterocycle is not more than 1. Examples of heterocycles include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl,
25 benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazoliny, carbazolyl, NH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2*H*,6*H*-1,5,2-dithiazinyl, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazoliny, imidazolyl, 1*H*-indazolyl, indolenyl, indolinyl, indoliziny, indolyl, 3*H*-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl,
30 isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, morpholinyl,

naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl;- 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, purinyl, 5 pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinolizinyl, quinoxalinyl, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4- 10 thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl.

Preferred heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, 15 pyrrolyl, pyrazolyl, pyrrolidinyl, imidazolyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazolyl, and isatinoyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

20 As used herein, the term "carbocyclic aryl" includes groups that contain 1 to 3 separate or fused rings and from 6 to about 18 ring atoms, without hetero atoms as ring members. Specifically preferred carbocyclic aryl groups include phenyl, and naphthyl including 1-naphthyl and 2-naphthyl.

25 The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio. As used herein, 30 "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds

wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the

5 conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic,

10 lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4, and the like. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by

15 conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th

20 ed., Mack Publishing Company, Easton, PA, p. 1418 (1985), the disclosure of which is hereby incorporated by reference.

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to formula I *in vivo* when such prodrug is administered

25 to a mammalian subject. Prodrugs of a compound of formula I are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of formula I wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug or compound of formula I is administered to a

30 mammalian subject, cleaves to form a free hydroxyl, free amino, or free sulfhydryl group,

respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formula I, and the like.

5 Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. A stable compound or stable structure is meant to imply a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an effective therapeutic agent. The term "therapeutically effective amount" of a compound of this invention means an
10 amount effective to antagonize abnormal level of CRF or treat the symptoms of affective disorder, anxiety or depression in a host.

 The compounds of general Formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing
15 conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may
20 be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups
25 or elixirs.

 Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of
30 sweetening agents, flavoring agents, coloring agents and preserving agents in order to

provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring

agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monoleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monoleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a

demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above.

5 The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland
10 fixed oil may be employed including synthetic mono-or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formula I may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared
15 by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

Compounds of general Formula I may be administered parenterally in a sterile
20 medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.

Typical subjects to which compounds of the invention may be administered will
25 be mammals, particularly primates, especially humans. For veterinary applications, a wide variety of subjects will be suitable, e.g. livestock such as cattle, sheep, goats, cows, swine and the like; poultry such as chickens, ducks, geese, turkeys, and the like; and domesticated animals particularly pets such as dogs and cats. For diagnostic or research applications, a wide variety of mammals will be suitable subjects including rodents (e.g.
30 mice, rats, hamsters), rabbits, primates, and swine such as inbred pigs and the like.

Additionally, for *in vitro* applications, such as *in vitro* diagnostic and research applications, body fluids and cell samples of the above subjects will be suitable for use such as mammalian, particularly primate such as human, blood, urine or tissue samples, or blood urine or tissue samples of the animals mentioned for veterinary applications.

5

Dosage levels of the order of from about 0.1 mg to about 140 mg of a compound of the invention per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1 mg to about 500 mg of an active ingredient.

10

Frequency of dosage may also vary depending on the compound used and the particular disease treated. However, for treatment of most CNS disorders, a dosage regimen of 4 times daily or less is preferred. For the treatment of stress and depression a dosage regimen of 1 or 2 times daily is particularly preferred.

15

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

20

Preferred compounds of the invention will have certain pharmacological properties. Such properties include, but are not limited to oral bioavailability, low toxicity, low serum protein binding and desirable *in vitro* and *in vivo* half-lives. Penetration of the blood brain barrier for compounds used to treat CNS disorders is necessary, while low brain levels of compounds used to treat peripheral disorders are often preferred.

25

30

Assays may be used to predict these desirable pharmacological properties. Assays used to predict bioavailability include transport across human intestinal cell monolayers, including Caco-2 cell monolayers. Toxicity to cultured hepatocytes may be used to predict compound toxicity. Penetration of the blood brain barrier of a compound in humans may be predicted from the brain levels of the compound in laboratory animals given the compound intravenously.

Serum protein binding may be predicted from albumin binding assays. Such assays are described in a review by Oravcová, et al. (Journal of Chromatography B (1996) volume 677, pages 1-27).

Compound half-life is inversely proportional to the frequency of dosage of a compound. *In vitro* half-lives of compounds may be predicted from assays of microsomal half-life as described by Kuhnz and Gieschen (Drug Metabolism and Disposition, (1998) volume 26, pages 1120-1127). Alternatively, compound half-life may be predicted from an *in vitro* microsomal assay such as the assay given in example 99b.

The present invention also pertains to methods for altering the activity of CRF receptors, said method comprising exposing cells expressing such receptors to an effective amount of a compound of the invention, wherein the compound is present in the solution at a concentration sufficient to specifically alter the signal transduction activity in response to CRF in cells expressing high levels of CRF1 receptors *in vitro*. This method includes altering the signal transduction activity of CRF receptors *in vivo*, e.g., in a patient given an amount of a compound of Formula I that would be sufficient to alter the signal transduction activity in response to CRF in cells expressing high levels of CRF1 *in vitro*. The amount of a compound that would be sufficient to alter the signal transduction activity in response to CRF receptors may be determined via an assay of

CRF receptor mediated signal transduction, such as an assay wherein the binding of CRF to a cell surface CRF receptor effects a changes in reporter gene expression.

5 The present invention also pertains to packaged pharmaceutical compositions for treating disorders responsive to C5a receptor modulation, e.g., eating disorders, depression or stress. The packaged pharmaceutical compositions include a container holding a therapeutically effective amount of at least one CRF1 receptor modulator as described supra and instructions for using the treating disorder responsive to CRF1 receptor modulation in the patient.

10

PREPARATION OF ARYLPYRAZINES

The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods described below, together with synthetic
15 methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include but are not limited to those methods described below. Each of the references cited below are hereby incorporated herein by reference. Preferred methods for the preparation of compounds of the present invention include, but are not limited to, those described in Scheme I, Scheme
20 II, and Scheme III. Those who are skilled in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention. All references cited herein are hereby incorporated in their entirety herein by reference. The following abbreviations are used herein:


25	AcOH	acetic acid
	DMF	N,N-dimethylformamide
	Et ₂ O	diethyl ether
	EtOAc	ethyl acetate
	EtOH	ethanol
30	NaH	sodium hydride

NaHMDS sodium bis(trimethylsilyl)amide

THF tetrahydrofuran

EX# example number

Scheme I (Method A)



5

According to the general method A, wherein R_1 and R_3 are as defined for formula I and Hal represents a halogen atom, suitably chloride or bromide, the halide in VI can be displaced by an amine or (thio)alkoxide nucleophile. Thus, aminopyrazine can be prepared from VI and an amine in the presence of a suitable transition metal catalyst such as but not limited to palladium(II) acetate or tris(dibenzylideneacetone)dipalladium(0), a ligand such as but not limited to 1,1'-bis(diphenylphosphine)ferrocene, 2,2'-bis(diphenylphosphine)-1,1'-binaphthyl, dicyclohexyl(2-biphenyl)phosphine, tricyclohexylphosphine, or tri-*tert*-butylphosphine, and a base such as sodium or potassium *tert*-butoxide in inert solvents such as but not limited to toluene, ethyleneglycol dimethyl ether, diglyme, DMF, or N-methylpyrrolidinone at temperatures ranging from ambient to 100°C. (Thio)alkoxypyrazines can be prepared by treating VI with a sodium or potassium salt of an alcohol or thiol in an inert solvent such as THF, DMF, N-methylpyrrolidinone, or methyl sulfoxide at ambient temperature or at elevated temperature up to the boiling point of the solvent employed. Halogenation may be accomplished by a variety of methods known in the art, including treatment with N-chlorosuccinimide, bromine, N-bromosuccinimide, pyridinium tribromide, triphenylphosphine dibromide, iodine, and N-iodosuccinimide in solvents such as but not limited to dichloromethane, acetic acid, or methyl sulfoxide. The bromopyrazine can be converted to arylpyrazine VII by a transition metal-catalyzed coupling reaction with a metalloaryl reagent ($Ar-[M]$). More commonly employed reagent/catalyst pairs include aryl boronic acid/palladium(0) (Suzuki reaction; N. Miyaura and A. Suzuki, Chemical Review 1995, 95, 2457), aryl trialkylstannane/palladium(0) (Stille reaction; T. N.

Mitchell, Synthesis 1992, 803), arylzinc/palladium(0) and aryl Grignard/nickel(II).


Palladium(0) represents a catalytic system made of a various combination of metal/ligand pair which includes, but not limited to, tetrakis(triphenylphosphine)palladium(0), palladium(II) acetate/tri(*o*-tolyl)phosphine, tris(dibenzylideneacetone)dipalladium(0)/tri-
 5 *tert*-butylphosphine and dichloro[1,1'-bis(diphenylphosphine)ferrocene]palladium(0).

Nickel(II) represents a nickel-containing catalyst such as [1,2-

bis(diphenylphosphino)ethane]dichloronickel(II) and [1,3-


bis(diphenylphosphino)propane]dichloronickel(II). The arylpyrazine VII, when X is NH, may be further transformed to VIII by N-alkylation. The N-H group is
 10 deprotonated by a strong base such as but not limited to alkali metal hydride, alkali metal amide, or alkali metal alkoxide in inert solvents such as but not limited to THF, DMF, or methyl sulfoxide. Alkylation may be conducted using alkyl halide, suitably bromide or iodide, at temperatures ranging from 0°C to 100°C.

Scheme II (Method B)




In an alternative way, compounds of Formula VII can be prepared as outlined in Scheme II. Transition metal-catalyzed coupling of the halo pyrazine VI as described in the Method A can provide the intermediate VIII. Oxidation of sterically less hindered
 20 nitrogen can be effected by using a variety of oxidizing agents known in the art, which includes *m*-chloroperoxybenzoic acid, trifluoroperacetic acid, hydrogen peroxide, and monoperoxyphthalic acid. The N-oxide can undergo rearrangement to give chloropyrazine IX upon the action of phosphorus oxychloride at temperatures ranging from ambient to 100°C. Displacement of the chloride with a nitrogen, oxygen, or sulfur
 25 nucleophile as described in the Method A can furnish the compounds of Formula VII.

Scheme III (Method C)



Yet another way of preparing compounds of formula VII is illustrated in the Scheme III. Compounds of formula X, 3,6-dialkyl-2,5-dichloropyrazines, can be prepared from 2-alkylglycine according to a known literature procedure (Ref: *Chemical and Pharmaceutical Bulletin of Japan* 1979, 27, 2027). Nucleophilic displacement of one chloride followed by Suzuki-type coupling at the other, as described in the Method A, can furnish the compounds of formula VII.

Scheme IV (Method D)



Still another way of preparing compounds of formula VII is illustrated in Scheme IV. Commercially available 2,6-Dichloropyrazine can undergo monosubstitution with nitrogen, oxygen, or sulfur nucleophile to give XI. Thus, X may react with an amine in solvents such as but not limited to dichloromethane, acetonitrile, THF, DMF, N-methylpyrrolidinone, methyl sulfoxide, methanol, ethanol, and isopropanol at temperatures ranging from 0°C to the boiling point of the solvent. In addition, X may

react with a sodium or potassium (thio)alkoxide in inert solvents such as but not limited to THF, DMF, N-methylpyrrolidinone, or methyl sulfoxide at temperatures ranging from 0°C to ambient temperature. Resulting monochloropyrazine **XI** can be halogenated by using the conditions described in the Method A to give a mixture of regioisomeric bromides **XIIa** and **XIIb**. Transition metal-catalyzed (hetero)aryl-aryl coupling of **XIIa**, as described in the Method A, followed by another halogenation can provide **VII** ($R_1 = \text{Hal}$, $R_3 = \text{Cl}$) which can be further converted to **VII** by displacing one or both of the halogen atoms, either sequentially or simultaneously, with a variety of nucleophiles (R_1 -[M] and R_3 -[M]), same or different, in the presence or absence of a transition metal catalyst. The aforementioned nucleophiles may include sodium or potassium (thio)alkoxide, alkylamine, and organometallic reagent such as but not limited to alkyl Grignard reagents, alkylboronic acids or its ester, or alkylstannanes. The aforementioned transition metal catalyst may represent palladium or nickel catalysts described in the Method A. The other regioisomeric bromide **XIIb** can also be converted to **VII** by changing the order of the transformation sequence. Those who are skilled in the art will also recognize that one can further change the order of transformations to prepare the compounds of Formula **VII** by way of the intermediate **XIII**.

As discussed above, preferred arylpyrazines of the invention exhibit good activity in standard *in vitro* CRF receptor binding assays, specifically the assay as specified in Example 96, which follows. References herein to "standard *in vitro* receptor binding assay" are intended to refer to that protocol as defined in Example 96 which follows. Generally preferred compounds preferred arylpyrimidines of the invention have an IC_{50} of about 10 micromolar or less, still more preferably and IC_{50} of about 100 nanomolar or less even more preferably an IC_{50} of about 10 nanomolar or less or even 1 nanomolar or less in such a defined standard *in vitro* CRF receptor binding assay as exemplified by Example 96 which follows.

EXAMPLES

The preparation of the compounds of the present invention is illustrated further by the following examples, which are not to be construed as limiting the invention in scope or spirit to the specific procedures and compounds described in them.

5 Commercial reagents were used without further purification. room or ambient temperature refers to 20 to 25°C. Concentration *in vacuo* implies the use of a rotary evaporator. TLC refers to thin layer chromatography. Proton nuclear magnetic resonance (¹H NMR) spectral data were obtained at 300 or 400 MHz. Mass spectral data were obtained either by CI or APCI methods.

10

Example 1

[N-(1-ethyl)propyl]-5-(2,4-dimethoxyphenyl)-3,6-dimethylpyrazine-2-amine

[Formula I: Ar=2,4-dimethoxyphenyl; R₂= -N-(1-ethyl)propyl; R₁=R₃=CH₃]


A. To a mixture of 2-chloro-3,6-dimethylpyrazine (0.83 mmol),
15 tris(dibenzylideneacetone)dipalladium(0) (2 mol%), and BINAP (6 mol%) in ethyleneglycol dimethyl ether (2 mL) under nitrogen is added 1-ethylpropylamine (1.0 mmol) followed by sodium tert-butoxide (1.25 mmol). The mixture is stirred at 70 – 80°C for 2.5 h, diluted with aqueous ammonium chloride, and extracted 1:1 hexane -Et₂O. Combined extracts are dried (sodium sulfate), filtered, concentrated, and
20 chromatographed on silica gel (10:1 to 4:1 hexane-EtOAc eluent) to give the aminopyrazine.

B. A solution of N-(1-ethyl)propyl-3,6-dimethylpyrazine-2-amine (0.72 g; 3.7 mmol) in dichloromethane (20 mL) is cooled to 0°C and N-bromosuccinimide (0.72 g; 4.1 mmol) is added in portions. After the addition, the mixture is further stirred for 1 h while being allowed to warm to room temperature. The mixture is then concentrated to a small volume in vacuo, triturated with hexane, filtered, washed with hexane, and the filtrate is concentrated and chromatographed on silica gel to give the bromide (1.07 g).

C. A mixed solution of 5-bromo-[N-(1-ethyl)propyl]-3,6-dimethylpyrazine-2-amine (0.40 g; 1.47 mmol) and tetrakis(triphenylphosphine)palladium(0) (33 mg; 2 mol%) in ethyleneglycol dimethyl ether (8 mL) is stirred at room temperature for 15 min, when 2,4-dimethoxybenzeneboronic acid (1.76 mmol) and an aqueous solution of sodium carbonate (1.0M, 4 mL) are added sequentially. The mixture is heated to 75 C with stirring for 1.5 h, then diluted with 0.1N sodium hydroxide and extracted twice with 1:1 hexane-ethyl ether. Combined extracts are dried (sodium sulfate), filtered, concentrated, and chromatographed on silica (4:1 to 1:1 hexane-EtOAc) to give the title compound (0.50 g): ¹H NMR (CDCl₃, 400 MHz) δ 0.95 (t, 6H), 1.6 (m, 4H), 2.2 (s, 3H), 2.4 (s, 3H), 3.75 (s, 3H), 3.85 (s, 3H), 3.95 (br d, 1H), 4.1 (br q, 1H), 6.5 (s, 1H), 6.55 (d, 1H) 7.2 (d, 1H); MS (CI) 330.


The Examples 2-20a in the Table I may be prepared following the methods described in Example 1.

Table I



EX#	RA	RB	Ar	¹ H NMR (CDCl ₃)	MS (CI)	NAME
2	H	1-ethylpropyl	2,4-dichlorophenyl	1.0 (t, 6H), 1.6 (m, 4H), 2.2 (s, 3H), 2.4 (s, 3H), 4.05 (br, 2H), 7.25 (d, 1H), 7.3 (d, 1H), 7.45 (s, 1H)	338	3,6-dimethyl-N-(1-ethylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
3	H	1-ethylpropyl	4-methoxy-2-methylphenyl	1.0 (t, 6H), 1.5 - 1.7 (m, 4H), 2.1 (s, 3H), 2.2 (s, 3H), 2.4 (s, 3H), 3.8 (s, 3H), 4.0 (br, 1H), 4.1 (br, 1H), 6.75 (d, 1H), 6.8 (s, 1H), 7.1 (s, 1H)	326	3,6-dimethyl-N-(1-ethylpropyl)-5-(2-methyl-4-methoxyphenyl)pyrazin-2-amine
4	H	1-ethylpropyl	3-cyanophenyl	0.95 (t, 6H), 1.6 (m, 4H), 2.4 (s, 3H), 2.45 (s, 3H), 4.1 (br, 2H), 7.5 (t, 1H), 7.6 (d, 1H), 7.75 (d, 1H), 7.85 (s, 1H)	295	3,6-dimethyl-N-(1-ethylpropyl)-5-(3-cyanophenyl)pyrazin-2-amine
5	H	1-methoxy-2-butyl	2,4-dichlorophenyl	1.0 (t, 3H), 1.7 (m, 2H), 2.2 (s, 3H), 2.4 (s, 3H), 3.4 (s, 3H), 3.55 (m, 2H), 4.25 (br, 1H), 4.55 (d, 1H), 7.25 (d, 1H), 7.3 (d, 1H), 7.45 (s, 1H)	354	3,6-dimethyl-N-(1-methoxy-2-butyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
6	H	1-methoxy-2-butyl	2,4-dimethoxyphenyl	1.0 (t, 3H), 1.7 (m, 2H), 2.2 (s, 3H), 2.4 (s, 3H), 3.4 (s, 3H), 3.55 (m, 2H), 3.75 (s, 3H), 3.85 (s, 3H), 4.25 (br, 1H), 4.4 (d, 1H), 6.5 (s, 1H), 6.6 (d, 1H), 7.2 (d, 1H)	346	3,6-dimethyl-N-(1-methoxy-2-butyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine

7	2-methoxymethyl-1-pyrrolidine	2,4-dichlorophenyl	1.9 (m, 4H), 2.2 (s, 3H), 2.6 (s, 3H), 3.3 (dd, 1H), 3.35 (s, 3H), 3.4 (br, 1H), 3.6 (dd, 1H), 3.8 (m, 1H), 4.6 (m, 1H), 7.3 (s, 2H), 7.5 (s, 1H)	366	2-(2,4-dichlorophenyl)-5-[2-(methoxymethyl)pyrrolidin-1-yl]-3,6-dimethylpyrazine
8	2-methoxymethyl-1-pyrrolidine	2,4-dimethoxyphenyl	1.9 (m, 4H), 2.2 (s, 3H), 2.6 (s, 3H), 3.3 (dd, 1H), 3.4 (s, 3H), 3.6 (dd, 1H), 3.75 (s, 3H), 3.85 (s, 3H), 4.6 (m, 1H), 6.5 (s, 1H), 6.6 (d, 1H), 7.2 (d, 1H)	358	2-(2,4-dimethoxyphenyl)-5-[2-(methoxymethyl)pyrrolidin-1-yl]-3,6-dimethylpyrazine
9	H	1-propylbutyl	1.0 (t, 6H), 1.3 - 1.6 (m, 8H), 2.2 (s, 3H), 2.35 (s, 3H), 4.0 (d, 1H), 4.25 (m, 1H), 7.25 (m, 2H), 7.45 (s, 1H)	366	3,6-dimethyl-N-(1-propylbutyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
10	H	1-propylbutyl	1.0 (t, 6H), 1.3 - 1.6 (m, 8H), 2.1 (s, 3H), 2.2 (s, 3H), 2.35 (s, 3H), 3.8 (s, 3H), 3.9 (d, 1H), 4.25 (m, 1H), 6.8 (m, 2H), 7.1 (d, 1H)	342	3,6-dimethyl-N-(1-propylbutyl)-5-(2-methyl-4-methoxyphenyl)pyrazin-2-amine
11	H	1-propylbutyl	1.0 (t, 6H), 1.3 - 1.6 (m, 8H), 2.15 (s, 3H), 2.2 (s, 3H), 2.35 (s, 3H), 4.0 (d, 1H), 4.25 (br, 1H), 7.1 (d, 1H), 7.2 (d, 1H), 7.25 (s, 1H)	346	3,6-dimethyl-N-(1-propylbutyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
12	H	1-propylbutyl	1.0 (t, 6H), 1.3 - 1.6 (m, 8H), 2.1 (s, 3H), 2.3 (s, 3H), 3.0 (s, 6H), 3.9 (d, 1H), 4.25 (m, 1H), 6.8 (d, 1H), 7.0 (s, 1H), 7.1 (d, 1H)	409	3,6-dimethyl-N-(1-propylbutyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl)pyrazin-2-amine
13	H	1-phenylpropyl	1.0 (t, 3H), 1.9 - 2.1 (m, 2H), 2.2 (s, 3H), 2.4 (s, 3H), 4.6 (d, 1H), 5.15 (q, 1H), 7.2 - 7.4 (m, 8H)	386	3,6-dimethyl-N-(3-phenylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
14	H	1-phenylpropyl	1.0 (t, 3H), 1.9 - 2.1 (m, 2H), 2.05 (s, 3H), 2.1 (s, 3H), 2.4 (s, 3H), 3.8 (s, 3H), 4.5 (d, 1H), 5.15 (q, 1H), 6.8 (m, 2H), 7.0 (d, 1H), 7.2 - 7.4 (m, 5H)	362	3,6-dimethyl-N-(3-phenylpropyl)-5-(2-methyl-4-methoxyphenyl)pyrazin-2-amine

15	H	1-phenylpropyl	4-chloro-2-methylphenyl	1.0 (t, 3H), 1.9 - 2.1 (m, 2H), 2.05 (s, 3H), 2.1 (s, 3H), 2.4 (s, 3H), 3.8 (s, 3H), 4.5 (d, 1H), 5.15 (q, 1H), 7.0 (d, 1H), 7.2 - 7.4 (m, 7H)	366	3,6-dimethyl-N-(3-phenylpropyl)-5-(2-chloro-4-methylphenyl)pyrazin-2-amine
16	H	1-phenylpropyl	4-dimethylamino-2-trifluoromethylphenyl	1.0 (t, 3H), 1.8 - 2.1 (m, 2H), 2.1 (s, 3H), 2.4 (s, 3H), 3.0 (s, 6H), 4.5 (d, 1H), 5.15 (q, 1H), 6.8 (d, 1H), 7.0 (s, 1H), 7.1 (d, 1H), 7.2 - 7.4 (m, 5H)	429	3,6-dimethyl-N-(3-phenylpropyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl)pyrazin-2-amine
17	H	1-propylbutyl	2,4-dimethoxyphenyl	1.0 (t, 6H), 1.3 - 1.6 (m, 8H), 2.2 (s, 3H), 2.35 (s, 3H), 3.75 (s, 3H), 3.85 (s, 3H), 3.9 (d, 1H), 4.25 (br, 1H), 6.5 (s, 1H), 6.55 (d, 1H), 7.2 (d, 1H)	358	3,6-dimethyl-N-(1-propylbutyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
18	H	1-phenylpropyl	2,4-dimethoxyphenyl	1.0 (t, 3H), 1.9 - 2.1 (m, 2H), 2.2 (s, 3H), 2.4 (s, 3H), 3.7 (s, 3H), 3.8 (s, 3H), 4.5 (d, 1H), 5.15 (q, 1H), 6.5 (s, 1H), 6.55 (d, 1H), 7.1 (d, 1H), 7.2 - 7.4 (m, 5H)	378	3,6-dimethyl-N-(3-phenylpropyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
19	H	1-ethylpropyl	4-chloro-2-methylphenyl	1.0 (t, 6H), 1.4 - 1.6 (m, 4H), 2.1 (s, 3H), 2.15 (s, 3H), 2.4 (s, 3H), 4.0 - 4.1 (br, 2H), 7.1 (d, 1H), 7.2 (d, 1H), 7.25 (s, 1H)	318	3,6-dimethyl-N-(1-ethylpropyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
20	H	1-ethylpropyl	4-dimethylamino-2-trifluoromethylphenyl	1.0 (t, 6H), 1.4 - 1.6 (m, 4H), 2.15 (s, 3H), 2.35 (s, 3H), 3.0 (s, 6H), 4.0 (d, 1H), 4.1 (m, 1H), 6.8 (d, 1H), 7.0 (s, 1H), 7.1 (d, 1H)	381	3,6-dimethyl-N-(1-ethylpropyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl)pyrazin-2-amine
20a	H		2,4-dichlorophenyl	0.89 (t, 3H), 1.6 (m, 2H), 2.14 (s, 3H), 2.42 (s, 3H), 3.44 (m, 2H), 4.16 (m, 1H), 7.43 (d, 1H), 7.50 (d, 1H), 7.72 (s, 1H); HCl salt	340,	2-{[5-(2,4-dimethylphenyl)-3,6-dimethylpyrazin-2-yl]amino}butan-1-ol

Example 21**[N-(1-ethyl)propyl]-3,6-dimethyl-5-(2,4,6-trimethylphenyl) pyrazine-2-amine**

[Formula I: Ar=2,4,6-trimethylphenyl; R₂= -N-(1-ethyl)propyl; R₁=R₃=CH₃]

5 To a solution of 5-bromo-[N-(1-ethyl)propyl]-3,6-dimethylpyrazine-2-amine (200 mg) obtained as in Example 1B in THF (4 mL) at room temperature is added [1,3-bis(diphenylphosphino)propane]dichloronickel(II) (40 mg). After 10 min, 2,4,6-trimethylphenylmagnesium bromide (1.0M in THF, 4 mL) is added slowly dropwise. The mixture is stirred at room temperature for 1 day, then refluxed overnight. The
10 resulting dark solution is poured into aqueous ammonium chloride and extracted twice with ether. Combined extracts are dried (sodium sulfate), filtered, concentrated, and chromatographed to give the desired product (87 mg): ¹H NMR (CDCl₃, 400 MHz) δ 1.0 (t, 6H), 1.5 – 1.7 (m, 4H), 1.95 (s, 3H), 2.1 (s, 3H), 2.3 (s, 3H), 2.35 (s, 3H), 3.95 (br, 1H), 4.1 (br, 1H), 6.9 (s, 2H).

15

Example 22**3-ethyl-[N-(1-ethyl)propyl]-6-methyl-5-(2,4,6-trimethylphenyl) pyrazine-2-amine**

[Formula I: Ar=2,4,6-trimethylphenyl; R₂= -N(1-ethyl)propyl; R₁= CH₂CH₃; R₃=CH₃]

To a solution of [N-(1-ethyl)propyl]-3,6-dimethyl-5-(2,4,6-
20 trimethylphenyl)pyrazine-2-amine (74 mg; 0.24 mmol) in THF (2 mL) at 0°C is added n-butyllithium (2.5M in hexane, 0.24 mL). After 10 min at 0°C, iodomethane (0.020 mL) is added. The mixture is stirred at 0°C for 10 min before being poured into aqueous ammonium chloride and extracted with Et₂O. The extract is dried (sodium sulfate), filtered, concentrated and the residue is purified by preparative TLC (10% EtOAc in
25 hexane, developed 3 times) to give the title compound (17 mg): ¹H NMR (CDCl₃, 400 MHz) δ 1.0 (t, 6H), 1.25 (t, 3H), 1.5 – 1.7 (m, 4H), 1.95 (s, 3H), 2.05 (s, 3H), 2.3 (s, 3H), 2.65 (q, 2H), 4.05 (m, 2H), 6.9 (s, 2H).


Example 23

3,6-diethyl-5-(2,4-dimethoxyphenyl)-[N-(1-ethylpropyl)]pyrazine-2-amine [Formula I: Ar=2,4-dimethoxyphenyl; R₂= -N(1-ethyl)propyl; R₁=R₃=CH₂CH₃]

2-Chloro-3,6-diethylpyrazine (*Chem. Pharm. Bull. Jap.*, 27, 2027 (1979)) is converted to the desired product following the procedures described in Example 1: ¹H NMR (CDCl₃, 300 MHz): δ 0.95 (t, 6H), 1.15 (t, 3H), 1.25 (t, 3H), 1.5 - 1.7 (m, 4H), 2.45 (q, 2H), 2.65 (q, 2H), 3.75 (s, 3H), 3.85 (s, 3H), 4.0 - 4.2 (br, 2H), 6.5 (s, 1H), 6.55 (d, 1H), 7.2 (d, 1H). LC-MS: 358

The Examples 24-48 in the Table II may be prepared following the general procedure described in Example 23.

Table II








EX #	RA	RB	Ar	¹ HNMR (CDCl ₃)	MS (CI)	Name
23a	H	1-ethylpropyl	2-methoxy-4-trifluoromethoxyphenyl	0.96 (t,6H), 1.15(t,3H), 1.28(t,3H), 1.57(m,2H), 1.67(m,2H), 2.42(m,2H), 2.66(q,2H), 3.78(s,3H), 4.10(br. 2H), 6.77(s,1H), 6.89(d,1H), 7.27(d,1H)	412	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxy-4-trifluoromethoxyphenyl)pyrazin-2-amine
23b	H	1-ethylpropyl	4-methoxy-2-trifluoromethoxyphenyl	0.96 (t,6H), 1.15(t,3H), 1.27(t,3H), 1.57(m,2H), 1.67(m,2H), 2.45(q,2H), 2.63(q,2H), 3.85(s,3H), 4.10(br. 2H), 6.86(brs,1H), 6.89(dd,1H), 7.29(d,1H)	412	3,6-diethyl-N-(1-ethylpropyl)-5-(2-trifluoromethoxy-4-methoxyphenyl)pyrazin-2-amine
23c	H	1-ethylpropyl	2-methoxy-4-trifluoromethoxyphenyl	0.96 (t,6H), 1.15(t,3H), 1.28(t,3H), 1.57(m,2H), 1.67(m,2H), 2.42(m,2H), 2.63(q,2H), 3.82(s,3H), 4.10(m, 2H), 7.14(s,1H), 7.28(d,1H), 7.38(d,1H)	396	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxy-4-trifluoromethoxyphenyl)pyrazin-2-amine





23d	H	1-ethylpropyl	4-methoxy-2-trifluoromethylphenyl	0.96 (t,6H), 1.11 (t,3H), 1.24 (t,3H), 1.57 (m,2H), 1.67 (m,2H), 2.34 (q,2H), 2.63 (q,2H), 3.87 (s,3H), 4.10 (m, 2H), 7.07 (dd,1H), 7.24 (d,1H), 7.25 (s,1H)	396	3,6-diethyl-N-(1-ethylpropyl)-5-(2-trifluoromethyl-4-methoxyphenyl)pyrazin-2-amine
23e	H	1-ethylpropyl	2-methoxy-4-methylphenyl	0.95 (t,6H), 1.14 (t,3H), 1.28 (t,3H), 1.55 (m,2H), 1.67 (m,2H), 2.38 (s,3H), 2.46 (m,2H), 2.64 (q,2H), 3.75 (s,3H), 4.10 (m, 2H), 6.74 (s,1H), 6.82 (d,1H), 7.14 (d,1H)	342	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxy-4-methylphenyl)pyrazin-2-amine
23f	H	1-ethylpropyl	2,4-difluoro-methoxyphenyl	0.96 (t,6H), 1.17 (t,3H), 1.27 (t,3H), 1.57 (m,2H), 1.69 (m,2H), 2.44 (m,2H), 2.64 (q,2H), 4.10 (m, 2H), 7.20 (s,1H), 7.22 (d,1H), 7.44 (d,1H)	466	3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-bis-trifluoromethoxyphenyl)pyrazin-2-amine
23g	H	1-ethylpropyl	2,4-dichlorophenyl	0.96 (t,6H), 1.14 (t,3H), 1.28 (t,3H), 1.57 (m,2H), 1.67 (m,2H), 2.43 (m,2H), 2.64 (q,2H), 4.12 (m, 2H), 7.25 (d,1H), 7.30 (dd,1H), 7.46 (d,1H)	366	3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
24	H	1-methoxy-2-butyl	2,4-dichlorophenyl		382	3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
25	H	1-methylpropyl	2,4-dichlorophenyl		352	3,6-diethyl-N-(1-methylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine

26	H	1-phenylethyl	2,4-dichlorophenyl		400	3,6-diethyl-N-(2-phenylethyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
27	H	1-propylbutyl	4-chloro-2-methylphenyl		374	3,6-diethyl-N-(1-propylbutyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
28	H	1-methoxy-2-butyl	4-chloro-2-methylphenyl		362	3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
29	H	1-methylpropyl	4-chloro-2-methylphenyl		332	3,6-diethyl-N-(1-methylpropyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
30	H	1-phenylethyl	4-chloro-2-methylphenyl		380	3,6-diethyl-N-(2-phenylethyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
31	H	1-propylbutyl	4-methoxy-2-methylphenyl		370	3,6-diethyl-N-(1-propylbutyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
32	H	1-methoxy-2-butyl	4-methoxy-2-methylphenyl		358	3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2-methyl-4-methoxyphenyl)pyrazin-2-amine
33	H	1-methylpropyl	4-methoxy-2-methylphenyl		328	3,6-diethyl-N-(1-methylpropyl)-5-(2-methyl-4-methoxyphenyl)pyrazin-2-amine





34	H	1-propylbutyl	2,4-dimethoxyphenyl		386	3,6-diethyl-N-(1-propylbutyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
35	H	1-methoxy-2-butyl	2,4-dimethoxyphenyl		374	3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
36	H	1-methylpropyl	2,4-dimethoxyphenyl		344	3,6-diethyl-N-(1-methylpropyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
37	H	1-phenylethyl	2,4-dimethoxyphenyl		392	3,6-diethyl-N-(2-phenylethyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
38	H	benzyl	2,4-dimethoxyphenyl		378	3,6-diethyl-N-benzyl-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
39	H	2-methyl-1-phenylpropyl	2,4-dimethoxyphenyl		420	
40	H	1-propylbutyl	2-chloro-4-(dimethylamino)phenyl		403	3,6-diethyl-N-(1-propylbutyl)-5-(2-chloro-4-dimethylaminophenyl)pyrazin-2-amine
41	H	1-methoxy-2-butyl	2-chloro-4-(dimethylamino)phenyl		391	3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2-chloro-4-dimethylaminophenyl)pyrazin-2-amine
42	H	1-methylpropyl	2-chloro-4-(dimethylamino)phenyl		361	3,6-diethyl-N-(1-methylpropyl)-5-(2-chloro-4-dimethylaminophenyl)pyrazin-2-amine





43	H	2-methylpropyl	2-chloro-4-(dimethylamino) phenyl		361	3,6-diethyl-N-(2-methylpropyl)-5-(2-chloro-4-dimethylaminophenyl)pyrazin-2-amine
44	H	1-phenylethyl	2-chloro-4-(dimethylamino) phenyl		409	3,6-diethyl-N-(2-phenylethyl)-5-(2-chloro-4-dimethylphenyl)pyrazin-2-amine
45	H	1-propylbutyl	4-dimethylamino-2-(trifluoromethyl) phenyl		437	3,6-diethyl-N-(1-propylbutyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl) pyrazin-2-amine
46	H	1-methoxy-2-butyl	4-dimethylamino-2-(trifluoromethyl) phenyl		425	3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl) pyrazin-2-amine
46a	H	1-ethylpropyl				5-(2-methoxy-4-methylpyrimidin-5-yl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
46b	H	1-ethylpropyl				5-(2-methyl-6-methoxypyridin-3-yl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
46c	H	1-ethylpropyl				5-(2-dimethylamino-6-methoxypyridin-3-yl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine


46d	H	1-ethylpropyl	5-chloro-2,4-dimethylphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-dimethyl-5-chlorophenyl)pyrazin-2-amine
46e	H	1-ethylpropyl	2,4-dichloro-5-methylphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-dimethyl-5-methylphenyl)pyrazin-2-amine
46f	H	1-ethylpropyl				5-[2,4-dimethyl-5-(2-morpholin-4-ylethoxy)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
46g	H	1-ethylpropyl	2,4-dimethyl-5-methoxyphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-dimethyl-5-methoxyphenyl)pyrazin-2-amine
46h	H	1-ethylpropyl	2-methyl-4-methoxyphenyl		342	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-methoxyphenyl)pyrazin-2-amine
46i	H	1-ethylpropyl	2-methyl-4-chlorophenyl		382	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
46j	H	1-ethylpropyl			345	5-[5-(1-ethylpropylamino)-3,6-dimethylpyrazin-2-yl]-6-methoxypyridin-2(3H)-one

46k	H	1-ethylpropyl			345	3-[5-(1-ethylpropylamino)-3,6-dimethylpyrazin-2-yl]-6-methoxypyridin-2(3H)-one
47	H	1-methylpropyl	4-dimethylamino-2-(trifluoromethyl)phenyl		395	3,6-diethyl-N-(1-methylpropyl)-5-(2-trifluoromethyl-4-methoxyphenyl)pyrazin-2-amine
47a	H		2-methoxy-4,6-dimethylphenyl	1.10 (t, 3H), 1.26 (t, 3H), 1.94 (s, 3H), 2.33 (s, 3H), 2.38 (m, 2H), 2.67 (m, 2H), 3.40 (s, 3H), 3.42 (s, 3H), 3.57 (m, 2H), 3.68 (m, 2H), 3.70 (s, 3H), 4.53 (m, 1H), 4.69 (m, 1H), 6.60 (s, 1H), 6.71 (s, 1H)	388	3,6-diethyl-N-[2-methoxy-1-(methoxymethyl)ethyl]-5-(2-methoxy-4,6-dimethylphenyl)pyrazin-2-amine
47b	H		2,4-dimethoxyphenyl	1.14 (t, 3H), 1.26 (t, 3H), 2.44 (m, 2H), 2.67 (m, 2H), 3.40 (s, 6H), 3.53 (m, 2H), 3.66 (m, 2H), 3.75 (s, 3H), 3.82 (s, 3H), 4.52 (m, 1H), 4.71 (m, 1H), 6.50 (s, 1H), 6.55 (d, 1H), 7.18 (d, 1H)	390	3,6-diethyl-N-[2-methoxy-1-(methoxymethyl)ethyl]-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
47c	H		2,4-dichlorophenyl	1.14 (t, 3H), 1.29 (t, 3H), 2.43 (b, 2H), 2.66 (m, 2H), 3.42 (s, 6H), 3.56 (m, 2H), 3.68 (m, 2H), 4.55 (m, 1H), 4.86 (d, 1H), 7.3 (m, 2H), 7.48 (m, 1H)	398	3,6-diethyl-N-[2-methoxy-1-(methoxymethyl)ethyl]-5-(2,4-dichlorophenyl)pyrazin-2-amine

47d	H		1.11 (t, 3H), 1.26 (t, 3H), 1.98 (s, 3H), 2.29 (m, 4H), 2.32 (s, 3H), 2.37 (m, 2H), 2.57 (m, 2H), 2.66 (m, 2H), 3.42 (s, 6H), 3.54 (m, 6H), 3.68 (m, 2H), 4.0 (m, 2H), 4.54 (m, 1H), 4.70 (d, 1H), 6.59 (s, 1H), 6.72 (s, 1H)	487	3,6-diethyl-N-[2-methoxy-1-(methoxymethyl)ethyl]-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazin-2-amine
47e	H		0.98 (t, 6H), 1.12 (t, 3H), 1.28 (t, 3H), 1.57 (m, 2H), 1.71 (m, 2H), 2.13 (s, 3H), 2.40 (m, 2H), 2.66 (m, 2H), 4.1 (b, 2H), 6.93 (m, 2H), 7.15 (m, 1H)	330	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-fluorophenyl)pyrazin-2-amine
47f	2-methoxyethyl		1.13 (t, 3H), 1.24 (t, 3H), 2.5 (b, 2H), 2.84 (q, 2H), 3.33 (s, 6H), 3.6 (m, 8H), 7.31 (m, 2H), 7.49 (s, 1H)	413	3,6-diethyl-N,N-(2-methoxyethyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
47g	Propyl		0.12 (m, 2H), 0.48 (m, 2H), 0.91 (t, 3H), 1.05 (m, 1H), 1.15 (m, 3H), 1.3 (m, 3H), 1.6 (m, 2H), 2.5 (b, 2H), 2.83 (m, 2H), 3.16 (m, 2H), 3.34 (m, 2H), 7.34 (s, 2H), 7.49 (s, 1H)	391	3,6-diethyl-N-propyl-N-(cyclopropylmethyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
47h	H		0.94 (t, 6H), 1.16 (t, 3H), 1.28 (t, 3H), 1.55 (m, 2H), 1.78 (m, 2H), 2.42 (q, 2H), 2.62 (q, 2H), 3.94 (s, 3H), 4.01 (s, 3H), 4.12 (m, 2H), 8.2 (s, 1H)	361	5-(2,4-dimethoxypyrimidin-5-yl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine

47i	H	1-ethylpropyl	2,4-bis(trifluoromethyl)-6-methoxyphenyl	0.98 (t, 6H), 1.12 (t, 3H), 1.25 (t, 3H), 2.64 (m, 4H), 2.29 (q, 2H), 2.61 (q, 2H), 3.81 (s, 3H), 4.12 (m, 2H), 7.33 (s, 1H), 7.60 (s, 1H)	464	3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-bis-trifluoromethyl-6-methoxyphenyl)pyrazin-2-amine
47j	H		2,4-dichlorophenyl	0.5 (m, 8H), 1.03 (m, 2H), 1.1 (t, 3H), 1.3 (t, 3H), 2.4 (b, 2H), 2.66 (m, 2H), 3.5 (m, 1H), 4.25 (d, 1H), 7.25 (m, 2H), 7.46 (s, 1H)	390	3,6-diethyl-N-(dicyclopropylmethyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
47k	H	NMe ₂ 	2,4-dichlorophenyl	1.16 (t, 3H), 1.28 (t, 3H), 1.35 (d, 3H), 2.25 (m, 2H), 2.29 (s, 6H), 2.5 (m, 2H), 2.68 (m, 2H), 4.06 (m, 1H), 5.15 (b, 1H), 7.28 (m, 2H), 7.47 (s, 1H)	381	N ² -[3,6-diethyl-(5-2,4-dichlorophenyl)pyrazin-2-yl]-N ¹ ,N ¹ -dimethylpropane-1,2-diamine
47l	H		2,4-dimethoxyphenyl	0.48 (m, 8H), 1.05 (m, 2H), 1.12 (t, 3H), 1.30 (t, 3H), 2.41 (q, 2H), 2.69 (q, 2H), 3.50 (m, 1H), 3.74 (s, 3H), 3.83 (s, 3H), 4.31 (d, 1H), 6.51 (s, 1H), 6.56 (d, 1H), 7.17 (d, 1H)	382	3,6-diethyl-N-(dicyclopropylmethyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
47m	H	1-ethylpropyl		0.97 (t, 6H), 1.15 (t, 3H), 1.28 (t, 3H), 1.6 (m, 4H), 2.46 (q, 2H), 2.65 (q, 2H), 3.91 (s, 3H), 3.94 (s, 3H), 4.07 (m, 2H), 6.40 (d, 1H), 7.50 (d, 1H)	359	3,6-diethyl-N-(1-ethylpropyl)-5-(2,6-dimethoxypyridin-3-yl)pyrazin-2-amine

47n	H			0.44 (m, 8H), 1.03 (m, 2H), 1.12 (t, 3H), 1.30 (t, 3H), 2.44 (q, 2H), 2.67 (q, 2H), 3.49 (m, 1H), 3.90 (s, 3H), 3.94 (s, 3H), 4.34 (d, 1H), 6.40 (d, 1H), 7.50 (d, 1H)	383	3,6-diethyl-N-(dicyclopropylmethyl)-5-(2,6-dimethoxypyridin-3-yl)pyrazin-2-amine
47o	Propyl	NMe ₂ 	2,4-dichlorophenyl	0.90 (t, 3H), 1.05 (t, 3H), 1.25 (t, 3H), 1.4 (m, 2H), 1.6 (m, 2H), 2.30 (s, 6H), 2.52 (m, 2H), 2.80 (m, 2H), 3.26 (m, 2H), 3.5 (b, 2H), 7.32 (m, 2H), 7.50 (s, 1H)	409	3,6-diethyl-N-propyl-N-(2-dimethylaminoethyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
47p	H	1-ethylpropyl	4-chloro-2-methylphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
47q	H	1-ethylpropyl	2-chloro-4-methoxyphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-methoxyphenyl)pyrazin-2-amine
47r	H	1-ethylpropyl				3,6-diethyl-N-(1-ethylpropyl)-5-(2-amino-4-methylpyridin-2-yl)pyrazin-2-amine
47s	Ethyl	1-ethylpropyl	2,4-dichlorophenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
47t	H	1-ethylpropyl	2-chloro-4-trifluoromethylphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-cyanophenyl)pyrazin-2-amine

47u	H	1-ethylpropyl	4-cyano-2-methylphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine
47v	H	1-ethylpropyl	4-cyano-2-trifluoromethylphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2-trifluoromethyl-4-chlorophenyl)pyrazin-2-amine
47w	H	1-ethylpropyl	2-cyano-4-methoxyphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2-cyano-4-methoxyphenyl)pyrazin-2-amine
47x	H	1-ethylpropyl				3,6-diethyl-N-(1-ethylpropyl)-5-(3 trifluoromethyl-5-chloropyridin-6-yl)pyrazin-2-amine
47y	H	1-ethylpropyl	2-cyano-4-trifluoromethylphenyl			3,6-diethyl-N-(1-ethylpropyl)-5-(2-cyano-4-trifluoromethylphenyl)pyrazin-2-amine
47z	Methyl	1-ethylpropyl	2-chloro-4-methoxyphenyl			3,6-diethyl-N-methyl-N-(1-ethylpropyl)-5-(2-chloro-4-methoxyphenyl)pyrazin-2-amine
48	H	1-phenylethyl	4-dimethylamino-2-(trifluoromethyl)phenyl	443		3,6-diethyl-N-(2-phenylethyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl)pyrazin-2-amine

Example 48a


3,6-diethyl-5-(2,4-diethoxy)phenyl-[N-(1-ethylpropyl)]pyrazine-2-amine [Formula I:
Ar=2,4-diethoxyphenyl; R₂= -N(1-ethyl)propyl; R₁=R₃=CH₂CH₃]

A: To a solution of 3,6-diethyl-5-(2,4-dimethoxyphenyl)-[N-(1-ethylpropyl)]pyrazine-2-amine (910mg, 2.54mmol) (obtained in example 23) in dichloromethane was added BBr₃ (1N,6ml) at -78°C. The mixture is stirred for 10 min then gradually warmed to room temperature and stirred for 3 hr before being poured into ice-water and extracted with dichloromethane. The aqueous layer was basified with sat. NaHCO₃ and extracted with dichloromethane. The combined extracts were dried (sodium sulfate), filtered, concentrated and the residue is purified by column (20% EtOAc in hexane) to give 3,6-diethyl-5-(2,4-dihydroxyphenyl)-[N-(1-ethylpropyl)]pyrazine-2-amine as a yellow oil (590 mg, 71%): ¹H NMR (CDCl₃, 400 MHz) δ 0.94 (t,6H), 1.30(t,3H), 1.36(t,3H), 1.57(m,2H), 1.67(m,2H), 2.62(q,2H), 2.86(q,2H), 4.15(m, 2H), 6.41(d,1H), 6.51(d,1H), 7.27(d,1H). LC-MS: 330 (M+1).

B: The above yellow oil (60mg, 0.182mmol) was dissolved in DMF (2ml) and alkylated with iodoethane (0.072 ml, 0.9 mmol) in the presence of K₂CO₃ (125 mg) at 75 °C for 2 hr. The reaction mixture was then diluted with water and extracted with EtOAc. The extracts were dried (sodium sulfate), filtered, concentrated and the residue is purified by column (2.5% MeOH in dichloromethane) to give the title compound as an oil (49 mg, 70%): ¹H NMR (CDCl₃, 400 MHz) δ 0.96 (t,6H), 1.13(t,3H), 1.27(m,6H), 1.42(t,3H), 1.57(m,2H), 1.67(m,2H), 2.47(m,2H), 2.66(q,2H), 3.95-4.15(m, 6H), 6.49(s,1H), 6.53(d,1H), 7.15(d,1H). LC-MS: 387 (M+1).

Examples 48b – 48k may be prepared according to the general procedure described in Example 48a.

Table IIa



EX#	R ₂	R ₄	¹ HNMR (CDCl ₃)	MS(Cl)	Name
48b	OH	OCH ₃	0.95 (t, 6H), 1.30 (t, 3H), 1.37 (t, 3H), 1.55 (m, 2H), 1.67 (m, 2H), 2.64 (q, 2H), 2.86 (q, 2H), 3.82 (s, 3H), 4.15 (m, 2H), 6.48 (dd, 1H), 6.59 (d, 1H), 7.31 (d, 1H)	344 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-hydroxy-4-methoxyphenyl)pyrazin-2-amine
48c	OH	OCH ₂ CH ₃	0.95 (t, 6H), 1.30 (t, 3H), 1.37 (t, 3H), 1.42 (t, 3H), 1.55 (m, 2H), 1.67 (m, 2H), 2.62 (q, 2H), 2.87 (q, 2H), 4.05 (q, 2H), 4.10 (m, 2H), 6.46 (dd, 1H), 6.56 (d, 1H), 7.30 (d, 1H)	358 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-hydroxy-4-ethoxyphenyl)pyrazin-2-amine
48d	OH	OCH(CH ₃) ₂	0.95 (t, 6H), 1.30 (t, 3H), 1.35 (d, 6H), 1.37 (t, 3H), 1.55 (m, 2H), 1.67 (m, 2H), 2.64 (q, 2H), 2.87 (q, 2H), 4.11 (m, 2H), 4.56 (m, 1H), 6.45 (dd, 1H), 6.56 (d, 1H), 7.28 (d, 1H)	372 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-[2-hydroxy-4-(2-propoxy)phenyl]pyrazin-2-amine
48e	OH	OCHF ₂	0.95 (t, 6H), 1.30 (t, 3H), 1.37 (t, 3H), 1.55 (m, 2H), 1.69 (m, 2H), 2.64 (q, 2H), 2.86 (q, 2H), 4.14 (m, 1H), 4.18 (brs, 1H), 6.54 (t, J=74Hz, 1H), 6.65 (d, 1H), 6.78 (s, 1H), 7.36 (d, 1H)	380 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-hydroxy-4-difluoromethoxyphenyl)pyrazin-2-amine

48f	OCH ₃	OH	0.95 (t,6H), 1.13 (t,3H), 1.29 (t,3H), 1.55 (m,2H), 1.67 (m,2H), 2.45 (q,2H), 2.70 (q,2H), 3.68 (s,3H), 4.10 (m, 2H), 6.32 (d,1H), 6.33 (s,1H), 6.99 (d,1H)	344 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxy-4-hydroxyphenyl)pyrazin-2-amine
48g	OCH ₃	OCH ₂ CH ₃	0.95 (t,6H), 1.14 (t,3H), 1.28 (t,3H), 1.43 (t,3H), 1.55 (m,2H), 1.67 (m,2H), 2.46 (q,2H), 2.65 (q,2H), 3.74 (s,3H), 4.07 (q, 2H), 4.10 (m,2H), 6.51 (d,1H), 6.54 (dd,1H), 7.15 (d,1H)	372 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxy-4-ethoxyphenyl)pyrazin-2-amine
48h	OCH ₃	OCH(CH ₃) ₂	0.95 (t,6H), 1.14 (t,3H), 1.28 (t,3H), 1.33 (d,6H), 1.55 (m,2H), 1.67 (m,2H), 2.46 (q,2H), 2.65 (q,2H), 3.74 (s,3H), 4.02 (d, 1H), 4.08 (m,1H), 4.59 (m,1H), 6.51 (d,1H), 6.54 (dd,1H), 7.15 (d,1H)	386 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-[2-methoxy-4-(2-propoxy)phenyl]pyrazin-2-amine
48i	OCH(CH ₃) ₂	OCH(CH ₃) ₂	0.95 (t,6H), 1.14 (t,3H), 1.18 (br. 6H), 1.28 (t,3H), 1.34 (d,6H), 1.55 (m,2H), 1.67 (m,2H), 2.46 (m,2H), 2.65 (q,2H), 4.02 (br, 1H), 4.10 (m,1H), 4.34 (m,1H), 4.55 (m,1H), 6.48 (d,1H), 6.54 (dd,1H), 7.15 (d,1H)	414 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-[2,4-bis-(2-propoxy)phenyl]pyrazin-2-amine
48j	OCHF ₂	OCHF ₂	0.96 (t,6H), 1.16 (t,3H), 1.28 (t,3H), 1.57 (m,2H), 1.69 (m,2H), 2.46 (q,2H), 2.63 (q,2H), 4.15 (m,2H), 6.51 (t, J=75Hz, 1H), 6.55 (t, J=74Hz, 1H), 7.03 (s,1H), 7.04 (d,1H), 7.35 (d,1H)	430 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2,4-bis-difluoromethoxyphenyl)pyrazin-2-amine
48k	OH	CH ₃	0.95 (t,6H), 1.14 (t,3H), 1.28 (t,3H), 1.55 (m,2H), 1.67 (m,2H), 2.38 (s,3H), 2.46 (q,2H), 2.65 (q,2H), 3.75 (s,3H), 4.02 (d, 1H), 4.08 (m,1H), 6.74 (s,1H), 6.69 (d,1H), 7.13 (d,1H)	342 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-hydroxy-4-methoxyphenyl)pyrazin-2-amine
48l	OCH ₂ CH ₂ OH	CH ₃	0.95 (t,6H), 1.16 (t,3H), 1.25 (t,3H), 1.55 (m,2H), 1.67 (m,2H), 2.38 (s,3H), 2.54 (q,2H), 2.68 (q,2H), 3.74 (t,2H), 4.11 (m, 2H), 4.22 (m,2H), 6.83 (d,1H), 6.87 (s,1H), 7.07 (d,1H)	372 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-hydroxyethoxy-4-methylphenyl)pyrazin-2-amine

						amine
48m	$\text{OCH}_2\text{CH}_2\text{OCH}_3$	CH_3		0.95 (t,6H), 1.12(t,3H), 1.25(t,3H), 1.55(m,2H), 1.67(m,2H), 2.36(s,3H), 2.48(m,2H), 2.63(q,2H), 3.23(s,3H), 3.58(t,2H), 4.02(d,1H), 4.04(t,2H), 4.11(m,1H), 6.76(s,1H), 6.83(d,1H), 7.14(d,1H)	386 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxyethoxy-4-methylphenyl)pyrazin-2-amine
48n	$\text{O}(\text{CH}_2)_2\text{N}(\text{CH}_3)_2$	CH_3		0.94 (t,6H), 1.12(t,3H), 1.25(t,3H), 1.55(m,2H), 1.67(m,2H), 2.15(s,6H), 2.36(s,3H), 2.48(m,2H), 2.58(t,2H), 2.63(q,2H), 4.00(t,2H), 4.02(d,1H), 4.11(m,1H), 6.74(s,1H), 6.83(d,1H), 7.14(d,1H)	399 (M+1)	3,6-diethyl-N-(1-ethylpropyl)-5-(2-dimethylaminoethoxy-4-methylphenyl)pyrazin-2-amine

Example 48o:**[N-(1-ethyl)propyl]-5-[(2-dimethylamino-4-methyl)pyridin-5-yl]-3,6-****diethylpyrazine-2-amine** [Formula I: Ar=(2-dimethylamino-4-methyl)pyridin-5-yl; R₂=5 -N(1-ethyl)propyl); R₁=R₃=CH₂CH₃]

5-Bromo-3,6-diethyl-[N-(1-ethyl)propyl]pyrazine-2-amine obtained in Example 23 reacts with [(2-dimethylamino-4-methyl)pyridin-5-yl]magnesium bromide as in Example 21 to give the title compound.

¹H NMR (CDCl₃, 400 MHz) δ 0.96 (t,6H), 1.14(t,3H), 1.28(t,3H), 1.57(m,2H),
10 1.67(m,2H),2.12(s,3H), 2.48(m,2H), 2.64(q,2H), 3.12(s,6H), 4.10(m, 2H), 6.43(s,1H),
7.89(s,1H). LC-MS: 356

Example 49**[N-(1-ethyl)propyl]-5-(2-methoxy-4,6-dimethylphenyl)-3,6-dimethylpyrazine-2-amine**15 [Formula I: Ar=2-methoxy-4,6-dimethylphenyl; R₂= -N(1-ethyl)propyl; R₁=R₃=CH₃]

A. To a solution of 2-chloro-3,6-dimethylpyrazine (0.71 g, 5.0 mmol) and tetrakis(triphenylphosphine)palladium(0) (140 mg, 2.5 mol%) in ethylene glycol dimethyl ether (30 mL) is added 2-methoxy-4,6-dimethylbenzeneboronic acid (1.08 g, 6.0 mmol), followed by addition of 1 M aqueous sodium carbonate solution (15 mL). The
20 mixture is stirred at 70 – 75°C overnight, allowed to cool, diluted by saturated aqueous sodium bicarbonate solution, and extracted twice with Et₂O. Combined extracts are dried (sodium sulfate), filtered, and concentrated *in vacuo*. The residue is purified by filtration through a short pad of silica gel to give 1.4g of crude product.

25 B. The crude material obtained above is dissolved in dichloromethane (20 mL), cooled to 0°C, and *m*-chloroperoxybenzoic acid (70%, 1.2g) is added in portions. After 4 hours at ambient temperature, the mixture is diluted by *n*-hexane (20 mL) and washed with 1 M aqueous sodium hydroxide solution. The organic phase separated is dried (sodium sulfate), filtered, concentrated *in vacuo*, and the residue is used directly in the
30 next step without further purification.

C. The crude N-oxide is dissolved in phosphorus oxychloride (10 mL) and the solution is heated at 80°C overnight. Evaporation of phosphorus oxychloride and usual aqueous work-up of the residue followed by chromatography on silica yields 2-aryl-5-chloro-3,6-dimethylpyrazine as a white solid (0.76g).

D. To a solution of the chloropyrazine obtained in C (260mg, 0.94 mmol) and tris(dibenzylideneacetone)dipalladium(0) (11mg) in toluene (10 mL) is added a 0.2 M solution of tri-*tert*-butylphosphine in toluene (0.10 mL). After 15 min at ambient temperature, 1-ethylpropylamine (0.14 mL) and potassium *t*-butoxide (1.0 M in THF, 1.4 mL) are added successively and the reaction mixture is heated at 80°C for 4 hours. The mixture is allowed to cool to room temperature, diluted by ether, washed with aqueous ammonium chloride solution, dried (sodium sulfate), filtered, concentrated, and chromatographed on silica to give the desired product as a white solid (250mg): ¹H NMR (CDCl₃, 400 MHz) δ 0.95 (t, 6H), 1.5 - 1.7 (m, 4H), 2.0 (s, 3H), 2.1 (s, 3H), 2.4 (s, 6H), 3.7 (s, 3H), 3.95 (br d, 1H), 4.1 (br q, 1H), 6.6 (s, 1H), 6.7 (s, 1H).

Examples 50 - 53 may be prepared according to the general procedure described in Example 49.

Example 50

N-benzyl-5-(2-methoxy-4,6-dimethylphenyl)-3,6-diethylpyrazine-2-amine [Formula I: Ar=2-methoxy-4,6-dimethylphenyl; R₂= -NHCH₂C₆H₅); R₁= R₃=CH₂CH₃] MS (CI) 376.

Example 51

5-(2-methoxy-4,6-dimethylphenyl)-3,6-diethylpyrazine-[N-(1-phenyl-2-methyl)propyl]-2-amine [Formula I: Ar=2-methoxy-4,6-dimethylphenyl; R₂= -NHCH(C₆H₅)CH(CH₃)CH₃; R₁= R₃=CH₂CH₃] MS (CI) 418.

Example 52**5-(2-methoxy-4,6-dimethylphenyl)-3,6-diethylpyrazine-[N-(1-propyl)butyl]-2-amine**

[Formula I: Ar=2-methoxy-4,6-dimethylphenyl; $R_2 = -NHCH(CH_2CH_2CH_3)_2$; $R_1 = R_3 = CH_2CH_3$] MS (CI) 384.

5

Example 53**[N-(1-methoxy)-2-butyl]-5-(2-methoxy-4,6-dimethylphenyl)-3,6-diethylpyrazine-2-**

amine [Formula I: Ar=2-methoxy-4,6-dimethylphenyl; $R_2 = -NCH(CH_2OCH_3)CH_2CH_3$; $R_1 = R_3 = CH_2CH_3$] MS (CI) 372.

10


Example 54**5-(2-methoxy-2,4-dimethylphenyl)-3,6-dimethyl-2-(3-pentyloxy)pyrazine** [Formula I: Ar=2-methoxy-4,6-dimethylphenyl; $R_2 = -OCH(CH_2CH_3)_2$; $R_1 = R_3 = CH_3$]

To an suspension of NaH (60% in mineral oil, 40mg) in DMF (0.5 mL) is added 3-pentanol (0.1 mL). The mixture is stirred at ambient temperature until hydrogen evolution ceased. To the resulting alkoxide solution is added an N-methylpyrrolidinone solution of 2-aryl-5-chloro-3,6-dimethylpyrazine (20mg in 0.5 mL solvent) obtained as in Example 49C. After 1 hour at room temperature, the mixture is heated to 70°C for another hour before it being allowed to cool and diluted with aqueous ammonium chloride solution and extracted twice with Et₂O. Combined organics are dried (sodium sulfate), filtered, concentrated, and chromatographed on silica gel to give the title compound as a colorless oil (15mg): ¹H NMR (CDCl₃, 400 MHz) δ 1.0 (t, 6H), 1.75 (m, 4H), 1.95 (s, 3H), 2.15 (s, 3H), 2.35 (s, 3H), 2.45 (s, 3H), 3.7 (s, 3H), 5.1 (quint, 1H), 6.6 (s, 1H), 6.7 (s, 1H); MS (CI) 329, 259.

25

Examples 55-62 in Table III may be prepared following the general procedure described in Examples 54 and 63.

Table III
















EX#	R	RB	Ar	¹ H NMR (CDCl ₃)	MS (CI)	NAME
55	Me		2-methoxy-4,6-dimethylphenyl	0.4 (m, 2H), 0.6 (m, 2H), 1.35 (m, 1H), 1.95 (s, 3H), 2.15 (s, 3H), 2.35 (s, 3H), 2.5 (s, 3H), 3.7 (s, 3H), 4.2 (d, 2H), 6.6 (s, 1H), 6.7 (s, 1H)	313, 259	3,6-dimethyl-2-(2-cyclopropylmethoxy)-5-(2-methoxyethoxy)-4,6-dimethylphenyl pyrazine
56	Me		2,4-dichlorophenyl	0.4 0.6 (m, 10H), 2.2 (s, 3H), 2.45 (s, 3H), 4.5 (t, 1H), 7.25 (d, 1H), 7.3 (d, 1H), 7.5 (s, 1H)	363, 269	3,6-dimethyl-2-(2-dicyclopropylmethoxy)-5-(2,4-dichlorophenyl) pyrazine
57	Me		2,4-dichlorophenyl		393, 269	3,6-dimethyl-2-(1-cyclohexylpropoxy)-5-(2,4-dichlorophenyl) pyrazine
58	Me		2,4-dichlorophenyl		381, 269	3,6-dimethyl-2-(1-cyclopentylpropoxy)-5-(2,4-dichlorophenyl) pyrazine
59	Me		2,4-dichlorophenyl		393, 269	3,6-dimethyl-2-(1-cyclobutylpropoxy)-5-(2,4-dichlorophenyl) pyrazine
60	Me		2,4-dichlorophenyl		399, 269	3,6-dimethyl-2-(cyclopropylphenylmethoxy)-5-(2,4-dichlorophenyl) pyrazine



61	Me		2,4-dichlorophenyl		393, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(2-propylcyclohexyl)oxy]pyrazine
61a	Et		4-fluoro-2-methylphenyl		331	3,6-dimethyl-2-(1-ethylpropoxy)-5-(2,4-dichlorophenyl)pyrazine
61b	Et		2,4-dichlorophenyl		399	2-[2-methoxy-1-(methoxymethyl)ethoxy]-3,6-diethyl-5-(2-methoxy-4-trifluoromethoxyphenyl)pyrazine
61b	Et		2-methoxy-4-trifluoromethoxyphenyl		413	3,6-diethyl-2-(1-ethylpropoxy)-5-(2-methoxy-4-trifluorophenyl)pyrazine
62	Et		2,4-dimethoxyphenyl	1.0 (t, 6H), 1.2 (t, 3H), 1.25 (t, 3H), 1.75 (quint, 4H), 2.5 (br, 2H), 2.8 (q, 2H), 3.75 (s, 3H), 3.85 (s, 3H), 5.1 (quint, 1H), 6.5 (s, 1H), 6.6 (d, 1H), 7.2 (d, 1H)	359, 289	3,6-diethyl-2-(1-ethylpropoxy)-5-(2,4-dimethoxyphenyl)pyrazine
62a	Me		2,4-dichlorophenyl	1.9 (m, 2H), 2.1 (m, 2H), 2.23 (s, 3H), 2.34 (s, 3H), 2.4 (m, 2H), 2.46 (s, 3H), 2.7 (b, 2H), 5.2 (b, 1H), 7.26 (dd, 1H), 7.33 (dd, 1H), 7.48 (d, 1H)	366, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(1-methylpiperidin-4-yl)oxy]pyrazine
62b	Me		2,4-dichlorophenyl		407, 269	3,6-dimethyl-2-(1-cyclohexylbutoxy)-5-(2,4-dichlorophenyl)pyrazine

62c	Me		2,4-dichlorophenyl			381, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(4-methoxycyclohexyl)oxy]pyrazine
62d	Me		2,4-dichlorophenyl	2.2 (m, 2H), 2.25 (s, 3H), 2.46 (s, 3H), 3.9-4.1 (m, 4H), 5.6 (m, 1H), 7.26 (d, 1H), 7.33 (d, 1H), 7.49 (s, 1H)		339, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-(tetrahydrofuran-3-yl)oxy]pyrazine
62e	Me		2,4-dichlorophenyl			253, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-(tetrahydro-2H-pyran-4-yl)oxy]pyrazine
62f	Me		2,4-dichlorophenyl	1.4-2.4 (m, 6H), 2.23 (s, 3H), 2.33 (s, 3H), 2.44 (s, 3H), 2.7 (m, 1H), 3.1 (m, 1H), 5.3 (m, 1H), 7.26 (d, 1H), 7.31 (d, 1H), 7.48 (s, 1H)		366, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(1-methylpiperidin-3-yl)oxy]pyrazine
62g	Me		2,4-dichlorophenyl			380, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(1-ethylpiperidin-3-yl)oxy]pyrazine
62h	Me		2,4-dichlorophenyl			366, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(1-aminopiperidin-4-yl)oxy]pyrazine

62i	Me		2,4-dichlorophenyl		352, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-(1-piperidin-4-yl)oxy pyrazine
62j	Me		2,4-dichlorophenyl		382, 269	2-{(3,6-dimethyl)-[5-(2,4-dichlorophenyl)pyrazin-2-yl]oxy}-N,N-diethylpropan-1-amine
62k	Me		2,4-dichlorophenyl		340, 269	3-{[5-(2,4-dichlorophenyl)-3,6-dimethylpyrazin-2-yl]oxy}butylamine
62l	Me		2,4-dichlorophenyl		352, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-(piperidin-3-yl)oxy pyrazine
62m	Me		2,4-dichlorophenyl	1.16 (t, 3H), 2.1 (m, 1H), 2.24 (s, 3H), 2.3-2.6 (m, 4H), 2.46 (s, 3H), 2.8 (m, 2H), 3.0 (m, 1H), 5.5 (m, 1H), 7.26 (d, 1H), 7.32 (d, 1H), 7.48 (s, 1H)	366, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(1-ethylpyrrolidin-3-yl)oxy]pyrazine
62n	Me		2,4-dichlorophenyl		352, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(1-methylpyrrolidin-3-yl)oxy]pyrazine

62o	Me		2,4-dichlorophenyl		338, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-[(pyrrolidin-3-yl)oxy]pyrazine
62p	Me		2,4-dichlorophenyl		383, 269	ethyl 3-{[5-(2,4-dichlorophenyl)-3,6-dimethylpyrazin-2-yl]oxy}butanoate
62q	Me		2,4-dichlorophenyl	1.14 (t, 3H), 1.26 (t, 3H), 2.05 (q, 2H), 2.19 (s, 3H), 2.54 (s, 3H), 4.22 (q, 2H), 5.12 (t, 1H), 7.25 (d, 1H), 7.33 (d, 1H), 7.48 (s, 1H)	383, 269	ethyl 2-{[5-(2,4-dichlorophenyl)-3,6-dimethylpyrazin-2-yl]oxy}propanoate
62r	Me		2,4-dichlorophenyl	2.44 (s, 3H), 2.64 (s, 3H), 6.53 (d, 2H), 7.32 (d, 1H), 7.43 (d, 1H), 7.56 (s, 1H), 7.70 (d, 2H)	346, 269	2,5-dimethyl-3-(2,4-dichlorophenyl)-6-(pyridin-4-yl)oxy]pyrazine
62s	Et		2-methoxy-4,6-dimethylphenyl	0.98 (t, 6H), 1.11 (t, 3H), 1.23 (t, 3H), 1.75 (m, 4H), 1.96 (s, 3H), 2.35 (s, 3H), 2.4 (m, 2H), 2.8 (m, 2H), 3.70 (s, 3H), 5.1 (m, 1H), 6.60 (s, 1H), 6.71 (s, 1H)	357, 287	3,6-diethyl-2-(1-ethylpropoxy)-5-(2-methoxy-4,6-dimethylphenyl)pyrazine
62t	Me		2,4-dichlorophenyl	0.97 (t, 6H), 1.74 (m, 4H), 2.24 (s, 3H), 2.44 (s, 3H), 5.11 (m, 1H), 7.28 (d, 1H), 7.32 (d, 1H), 7.48 (s, 1H)	340, 269	3,6-dimethyl-2-(1-ethylpropoxy)-5-(2,4-dichlorophenyl)pyrazine

62u	Et			0.98 (m, 6H), 1.12 (t, 3H), 1.21 (t, 3H), 2.74 (m, 4H), 2.01 (s, 3H), 2.29 (s, 3H), 2.5 (m, 2H), 2.73 (m, 2H), 5.14 (m, 1H), 6.4 (b, 1H), 6.58 (s, 1H), 6.65 (s, 1H)	343, 273	3,6-diethyl-2-(1-ethylpropoxy)-5-(2-hydroxy-4,6-dimethylphenyl)pyrazine
62v	Et		2,4-dichlorophenyl	0.99 (t, 6H), 1.16 (t, 3H), 1.24 (t, 3H), 1.75 (m, 4H), 2.5 (b, 2H), 2.82 (m, 2H), 5.13 (m, 1H), 7.26 (d, 1H), 7.32 (d, 1H), 7.49 (s, 1H)	367, 297	3,6-diethyl-2-(1-ethylpropoxy)-5-(2,4-dichlorophenyl)pyrazine
62w	Et			0.99 (t, 6H), 1.15 (t, 3H), 1.23 (t, 3H), 1.72 (m, 4H), 1.99 (s, 3H), 2.25 (m, 4H), 2.32 (s, 3H), 2.4 (m, 2H), 2.55 (m, 2H), 2.8 (m, 2H), 2.56 (m, 4H), 4.0 (m, 2H), 5.14 (m, 1H), 6.57 (s, 1H), 6.71 (s, 1H)	456, 386	3,6-diethyl-2-(1-ethylpropoxy)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine
62x	Et		2,4-bis(trifluoromethyl)phenyl			3,6-diethyl-2-(1-ethylpropoxy)-5-(2,4-bis-trifluoromethylphenyl)pyrazine
62y	Et		2-chloro-4-trifluoromethylphenyl			3,6-diethyl-2-(1-ethylpropoxy)-5-(2-chloro-4-trifluorophenyl)pyrazine

62z	Et					3,6-diethyl-2-(1-ethylpropoxy)- 5-(2-methoxy-4-methylpyridin-3- yl)pyrazine
-----	----	---	---	--	--	---

Example 63


[N-(1-ethyl)propyl]-3,6-dimethyl-5-{2-[2-(4-morpholino)ethyl]oxy-4,6-dimethylphenyl}pyrazine-2-amine [Formula I: Ar=2-[2-(4-morpholino)ethyl]oxy-4,6-dimethylphenyl; R₂= -NHCH(CH₂CH₃)₂; R₁= R₃=CH₃]

- 5 A. A solution of 2-chloro-3,6-dimethyl-5-(2-methoxy-4,6-dimethylphenyl)pyrazine (180mg) in dichloromethane was cooled to 0°C and boron tribromide (0.1 mL) is added slowly dropwise. After the addition, the mixture is further stirred at 0°C for 1.5 hours, diluted by Et₂O, washed with saturated aqueous sodium bicarbonate solution, dried (sodium sulfate), filtered, and concentrated *in vacuo*. The
- 10 residue is used in the next step without further purification.
- B. To a suspension of the crude phenol and potassium carbonate (400mg) in DMF (4 mL) is added 4-(2-chloroethyl)morpholine hydrochloride (200mg) in one portion and the mixture is stirred at 60°C for 4 hours. After further stirring at room temperature overnight, the mixture is poured into aqueous sodium bicarbonate and extracted twice
- 15 with Et₂O - hexane. Combined extracts are dried (sodium sulfate), filtered, concentrated, and chromatographed on silica (eluent 5% triethylamine in 1:1 EtOAc - hexane) to give the product as a colorless oil (160mg).
- C. The chloropyrazine is converted to the corresponding aminopyrazine
- 20 following the same procedure as in Example 49D: ¹H NMR (CDCl₃, 400 MHz) δ 0.95 (m, 6H), 1.6 (m, 4H), 2.0 (s, 3H), 2.1 (s, 3H), 2.3 (m, 4H), 2.35 (s, 6H), 2.6 (m, 2H), 3.6 (m, 2H), 3.9 (d, 1H), 4.0 (m, 2H), 4.05 (m, 1H), 6.55 (s, 1H), 6.7 (s, 1H); MS (CI) 427.

Examples 64-74 in Table IV may be prepared following the general procedure described in Example 63.

Table IV

EX#	R	R _A	R _B	¹ H NMR (CDCl ₃)	MS (CI)	NAME
64	Me	H	1-propylbutyl	0.95 (t, 6H), 1.3 - 1.6 (m, 8H), 2.0 (s, 3H), 2.1 (s, 3H), 2.3 (m, 10H), 2.6 (m, 2H), 3.6 (m, 4H), 3.85 (d, 1H), 4.0 (m, 2H), 4.25 (m, 1H), 6.6 (s, 1H), 6.7 (s, 1H)	455	3,6-dimethyl-N-(1-propylbutyl)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
65	Me	H	1-methoxy-2-butyl	1.0 (t, 2H), 1.8 (m, 2H), 2.0 (s, 3H), 2.1 (s, 3H), 2.3 (m, 10H), 2.6 (br, 2H), 3.4 (s, 3H), 3.5 - 3.6 (m, 6H), 4.0 (br, 2H), 4.25 (br, 1H), 4.4 (d, 1H), 6.6 (s, 1H), 6.7 (s, 1H)	443	3,6-dimethyl-N-(1-methoxy-2-butyl)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
66	Me	propyl	propyl	0.9 (t, 6H), 1.6 (q, 4H), 2.0 (s, 3H), 2.15 (s, 3H), 2.3 (m, 4H), 2.35 (s, 3H), 2.45 (s, 3H), 2.55 (m, 2H), 3.2 (m, 4H), 3.6 (m, 4H), 4.0 (m, 2H), 6.6 (s, 1H), 6.7 (s, 1H)	441	3,6-dimethyl-N,N-dipropyl-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine



67	Et	H	1-ethylpropyl	0.95 (m, 6H), 1.1 (t, 3H), 1.25 (t, 3H), 1.5 - 1.7 (m, 4H), 2.0 (s, 3H), 2.25 (m, 4H), 2.3 (s, 3H), 2.35 (q, 2H), 2.55 (m, 2H), 2.6 (q, 2H), 3.55 (m, 4H), 4.0 (m, 3H), 4.1 (m, 1H), 6.6 (s, 1H), 6.7 (s, 1H)	455	3,6-diethyl-N-(1-ethylpropyl)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
68	Et	H	1-propylbutyl	0.95 (t, 6H), 1.1 (t, 3H), 1.2 (t, 3H), 1.4 (br, 4H), 1.55 (br, 4H), 2.0 (s, 3H), 2.25 (m, 4H), 2.3 (s, 3H), 2.35 (q, 2H), 2.55 (br s, 2H), 2.6 (q, 2H), 3.6 (br s, 4H), 4.0 (br, 3H), 4.25 (br q, 1H), 6.6 (s, 1H), 6.7 (s, 1H)	483	3,6-diethyl-N-(1-propylbutyl)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
69	Et	propyl	propyl	0.9 (t, 6H), 1.1 (t, 3H), 1.3 (t, 3H), 1.6 (m, 4H), 2.0 (s, 3H), 2.25 (br, 4H), 2.3 (s, 3H), 2.4 (m, 2H), 2.55 (br s, 2H), 2.7 - 2.9 (m, 2H), 3.2 (m, 4H), 3.6 (m, 4H), 4.0 (m, 2H), 6.6 (s, 3H), 6.7 (s, 1H)	469	3,6-diethyl-N,N-dipropyl-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
70	Et	ethyl	butyl	0.9 (t, 3H), 1.1 - 1.3 (m, 9H), 1.3 (m, 2H), 1.55 (m, 2H), 2.0 (s, 3H), 2.25 (br, 4H), 2.3 (s, 3H), 2.4 (m, 2H), 2.55 (br s, 2H), 2.7 - 2.9 (m, 2H), 3.2 (m, 4H), 3.6 (m, 4H), 4.0 (m, 2H), 6.6 (s, 3H), 6.7 (s, 1H)	469	3,6-diethyl-N-ethyl-N-propyl-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
71	Et	H	1-methylpropyl	1.0 (t, 3H), 1.1 (t, 3H), 1.2 (t + d, 6H), 1.5 - 1.7 (m, 2H), 2.0 (s, 3H), 2.25 (br s, 4H), 2.3 (s, 3H), 2.35 (q, 2H), 2.55 (br s, 2H), 2.6 (q, 2H), 3.55 (app t, 4H), 4.0 (m, 3H), 4.2 (m, 1H), 6.6 (s, 1H), 6.7 (s, 1H)	441	3,6-diethyl-N-(1-methylpropyl)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
72	Et	H	1-methoxy-2-butyl	1.0 (t, 3H), 1.1 (t, 3H), 1.3 (t, 3H), 1.7 (br, 2H), 2.0 (s, 3H), 2.25 (br s, 4H), 2.3 (s, 3H), 2.35 (q, 2H), 2.55 (br s, 2H), 2.6 (q, 2H), 3.4 (s, 3H), 3.6 (br s, 6H), 4.0 (br, 2H), 4.25 (br, 1H), 4.45 (d, 1H), 6.6 (s, 1H), 6.7 (s, 1H)	471	3,6-diethyl-N-(1-methoxy-2-butyl)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine

73	Et	propyl	cyclopropane- methyl	0.1 (app d, 2H), 0.45 (app d, 2H), 0.9 (t, 3H), 1.05 (m, 1H), 1.1 (t, 3H), 1.2 (t, 3H), 2.0 (s, 3H), 2.25 (m, 4H), 2.3 (s, 3H), 2.4 (m, 2H), 2.55 (br s, 2H), 2.8 (m, 2H), 3.1 (d, 2H), 3.3 (m, 2H), 3.6 (br s, 4 H), 4.0 (br, 2H), 6.6 (s, 1H), 6.7 (s, 1H)	481	3,6-diethyl-N-(1-methoxy-2-butyl)-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine
74	Et	ethyl	benzyl	1.0 - 1.3 (m, 9H), 2.0(s, 3H), 2.25 (br s, 4H), 2.3 (s, 3H), 2.4 (m, 2H), 2.55 (br s, 2H), 2.8 (m, 2H), 3.25 (q, 2H), 3.6 (br s, 4H), 4.0 (br, 2H), 4.45 (app q, 2H), 6.6 (s, 1H), 6.7 (s, 1H), 7.25 (m, 1H), 7.3 (t, 2H), 7.4 (d, 2H)	503	3,6-diethyl-N-ethyl-N-benzyl-5-[2,4-dimethyl-6-(2-morpholin-4-ylethoxy)phenyl]pyrazine-2-amine

Example 75**3-bromo-6-chloro-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]pyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; R₁=Br; R₂=-NHCH(CH₂CH₃)₂; R₃=Cl]

5 A. A solution of 2,6-dichloropyrazine (2.2g) and 1-ethylpropylamine (5 mL) in EtOH (10 mL) is heated at 140°C in a Teflon-sealed tube. Resulting solution is concentrated *in vacuo*, diluted by water, and extracted twice with hexane-ethyl ether. Combined extracts are dried (sodium sulfate), filtered, concentrated *in vacuo*, and the residue is filtered through a short pad of silica gel. The filtrate is concentrated to give 2-(3-pentylamino)-6-chloropyrazine as brownish oil that solidifies on standing (3.0g).

10

B. A solution of the above amine (4.09 g; 20.48 mmol) in chloroform (80 ml) is cooled to 0°C and N-bromosuccinimide (3.65 g; 20.48 mmol) is added in portions. The mixture is stirred at 0°C for 30 min, poured into saturated aqueous NaHCO₃, and extracted with dichloromethane. The combined extracts are washed successively with water and brine, dried (sodium sulfate), and concentrated *in vacuo*. The residue is chromatographed on silica gel (6% EtOAc in hexane) to give the desired 3-bromopyrazine as a minor product (0.53g; 9%) along with 6-chloro-5-bromo-[N-(1-ethyl)propyl]pyrazine-2-amine (4.37 g; 77%) as the major isomer.

15

20 C. 5-bromopyrazine obtained as above undergoes Suzuki coupling with 2,4-dichlorobenzeneboronic acid following the procedures in Example 1C to give 6-chloro-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]pyrazine-2-amine.

D. The aryl pyrazine is brominated again by using N-bromosuccinimide as described in Example 75B to give the desired product: ¹H NMR (CDCl₃, 400 MHz) δ 0.97 (t, 6H), 1.58 (m, 2H), 1.70 (m, 2H), 4.00 (m, 1H), 5.20 (d, 1H), 7.30 (s, 2H), 7.50 (s, 1H); MS(CI) 422.

25

Example 76

6-chloro-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]-3-(2-propenyl)pyrazine-2-amine

[Formula I: Ar=2,4-dichlorophenyl; R₁=CH₂CH=CH₂; R₂= -NHCH(CH₂CH₃)₂; R₃=Cl]

The bromopyrazine obtained in Example 75 is converted to the desired product by using allylboronic acid following the same procedure as in Example 1C: ¹H NMR

5 (CDCl₃, 400 MHz) δ 0.93 (t, 6H), 1.50 – 1.71 (m, 4H), 3.50 (d, 2H), 4.05 (m, 1H), 4.55 (d, 1H), 5.22 (m, 2H), 5.92 (m, 1H), 7.34 (m, 2H), 7.48 (m, 1H); MS(CI) 384.

Example 77**6-chloro-5-(2,4-dichlorophenyl)-3-ethyl-[N-(1-ethyl)propyl]pyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; R₁=CH₂CH₃; R₂= -NHCH(CH₂CH₃)₂; R₃=Cl]

10 The bromopyrazine obtained in Example 75 is converted to the desired product by using ethaneboronic acid following the same procedure as in Example 1C: ¹H NMR (CDCl₃, 400 MHz) δ 0.96 (t, 6H), 1.30 (t, 3H), 1.56 (m, 2H), 1.70 (m, 2H), 2.65 (t, 2H), 4.08 (m, 1H), 4.35 (d, 1H), 7.32 (d, 1H), 7.33 (s, 1H), 7.48 (d, 1H); MS (CI) 372.

15 Example 78

5-(2,4-dichlorophenyl)-6-ethyl-[N-(1-ethyl)propyl]pyrazine-2-amine [Formula I: Ar=2,4-dichlorophenyl; R₁=H; R₂= -NHCH(CH₂CH₃)₂; R₃=CH₂CH₃]

A. 6-chloro-[N-(1-ethyl)propyl]pyrazine-2-amine obtained in Example 75A reacts with ethylmagnesium bromide as in Example 21 to give 6-ethyl-[N-(1-ethyl)propyl]pyrazine-2-amine.

25 B. The above amine is brominated and coupled with 2,4-dichlorobenzeneboronic acid following the same procedure as described in Examples 75B and 75C, respectively to give the title compound: ¹H NMR (CDCl₃, 400 MHz) δ 0.97 (t, 6H), 1.13 (t, 3H), 1.56 (m, 2H), 1.65 (m, 2H), 2.45 (m, 2H), 3.72 (m, 1H), 4.45 (d, 1H), 7.25 (d, 1H), 7.30 (dd, 1H), 7.48 (d, 1H), 7.74 (s, 1H); MS(CI) 338.


Example 79**3-bromo-5-(2,4-dichlorophenyl)-6-ethyl-[N-(1-ethyl)propyl]pyrazine-2-amine**

30 [Formula I: Ar=2,4-dichlorophenyl; R₁=Br; R₂= -NHCH(CH₂CH₃)₂; R₃=CH₂CH₃]

The product of Example 78 is brominated as in Example 75B to give the title compound: ^1H NMR (CDCl_3 , 400 MHz) δ 0.97 (t, 6H), 1.14 (t, 3H), 1.56 (m, 2H), 1.65 (m, 2H), 2.45 (m, 2H), 4.02 (m, 1H), 5.00 (d, 1H), 7.25 (d, 1H), 7.30 (dd, 1H), 7.46 (d, 1H); MS (CI) 416.

Examples 80-82 in Table V may be prepared following the general procedure described in Example 79 using the corresponding halogenating agent indicated in the table.

Table V					
EX#	X	Reagent	¹ H NMR	MS(Cl)	NAME
80	F	SelectFluor(TM)	0.96 (t, 6H), 1.13 (t, 3H), 1.50 - 1.72 (m, 4H), 2.42 (m, 2H), 4.05 (m, 1H), 4.60 (d, 1H), 7.24 (d, 1H), 7.30 (dd, 1H), 7.48 (d, 1H).	356	3-fluoro-6-ethyl-N-(1-ethylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
81	Cl	N-chlorosuccinimide	0.97 (t, 6H), 1.14 (t, 3H), 1.55 (m, 2H), 1.70 (m, 2H), 2.46 (m, 2H), 4.05 (m, 1H), 4.93 (d, 1H), 7.24 (d, 1H), 7.30 (dd, 1H), 7.47 (d, 1H).	372	3-chloro-6-ethyl-N-(1-ethylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine
82	I	N-iodosuccinimide	0.97 (t, 6H), 1.14 (t, 3H), 1.55 (m, 2H), 1.68 (m, 2H), 2.43 (m, 2H), 4.00 (m, 1H), 4.95 (d, 1H), 7.24 (d, 1H), 7.30 (dd, 1H), 7.46 (d, 1H).	464	3-iodo-6-ethyl-N-(1-ethylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine



Example 83**5-(2,4-dichlorophenyl)-6-ethyl-[N-(1-ethyl)propyl]-3-methoxypyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; R₁=OCH₃; R₂= -NHCH(CH₂CH₃)₂; R₃=CH₂CH₃]

To a 1 N solution of sodium methoxide in N-methylpyrrolidinone is added 3-bromo-6-ethyl-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl] pyrazine-2-amine obtained as in Example 79. The mixture is heated to 70°C for 6 hr before being allowed to cool and diluted with water and the reaction mixture is extracted with 20% EtOAc in hexane. The combined extracts are washed with water, dried (sodium sulfate), filtered, concentrated *in vacuo*, and chromatographed on silica gel to give the title compound: ¹H NMR (CDCl₃, 400 MHz) δ 0.97 (t, 6H), 1.12 (t, 3H), 1.56 (m, 2H), 1.65 (m, 2H), 2.40 (q, 2H), 3.92 (s, 3H), 4.04 (m, 1H), 4.82 (d, 1H), 7.28 (m, 2H), 7.48 (d, 1H); MS(Cl) 368.

Example 84**5-(2,4-dichlorophenyl)-3-ethyl-6-methyl-[N-(1-ethyl)propyl]pyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; R₁=CH₂CH₃; R₂= -NHCH(CH₂CH₃)₂; R₃=CH₃]

A: To a solution of 2-(3-pentylamino)-6-chloropyrazine (4.26g, 21.3mmol) (obtained in example 75) in THF (30 mL) at room temperature is added [1,3-bis(diphenylphosphino)propane]dichloronickel(II) (540 mg). After 10 min, methylmagnesium bromide (3.0M in diethyl ether, 15.7 mL) is added slowly dropwise at 0 °C. The mixture is stirred at room temperature for 1 hr. The resulting dark solution is poured into aqueous ammonium chloride and extracted twice with ether. Combined extracts are dried (sodium sulfate), filtered, concentrated, and chromatographed to give the desired product as a light brown oil(98%).

B: A solution of the above oil in chloroform(60 mL) is cooled to 0°C and N-bromosuccinimide (3.8 g) is added in portions. After the addition, the mixture is further stirred for 1 h while being allowed to warm to room temperature. The mixture is then concentrated to a small volume in *vacuo*, triturated with hexane, filtered, washed with hexane, and the filtrate is concentrated and chromatographed on silica gel (8% ethyl

acetate in hexane elution) to give 5-bromo-[N-(1-ethyl)propyl]-6-methylpyrazine-2-amine (92%).

C. A mixed solution of the above bromide (1.2 g; 4.65 mmol) and
5 tetrakis(triphenylphosphine)palladium(0) (4 mol%) in ethyleneglycol dimethyl ether (60 mL) is stirred at room temperature for 15 min, when 2,4-dichlorobenzeneboronic acid (1.3g, 7 mmol) and an aqueous solution of sodium carbonate (1.0M, 12 mL) are added sequentially. The mixture is heated to 75°C with stirring for 1.5 h, then diluted with
10 0.1N sodium hydroxide and extracted twice with 1:1 hexane-ethyl ether. Combined extracts are dried (sodium sulfate), filtered, concentrated, and chromatographed on silica (4:1 hexane-EtOAc) to give 5-(2,4-dichlorophenyl)-6-methyl-[N-(1-ethyl)propyl]pyrazine-2-amine as a yellow oil (1.46 g, 97%): ¹H NMR (CDCl₃, 400 MHz) δ 0.97 (t,6H), 1.54(m,2H), 1.67(m,2H), 2.22(s,3H), 3.65(m,1H), 4.50(br, 1H), 7.27(d,1H), 7.31(dd,1H), 7.48(d,1H), 7.76(s,1H).

15

D: A solution of the above oil (1.27g, 3.92 mmol) in chloroform(40 mL) is cooled to 0°C and N-bromosuccinimide (770 mg) is added in portions. After the addition, the mixture is further stirred for 1 h while being allowed to warm to room temperature. The mixture is then concentrated to a small volume in vacuo, triturated
20 with hexane, filtered, washed with hexane, and the filtrate is concentrated and chromatographed on silica gel (3% ethyl acetate in hexane elution) to give 3-bromo-5-(2,4-dichlorophenyl)-6-methyl-[N-(1-ethyl)propyl]pyrazine-2-amine (1.56g, 98%).

E: A mixed solution of 3-bromo-5-(2,4-dichlorophenyl)-6-methyl-[N-(1-ethyl)propyl]pyrazine-2-amine (960 mg; 2.38 mmol) and
25 tetrakis(triphenylphosphine)palladium(0) (4 mol%) in ethyleneglycol dimethyl ether (25 mL) is stirred at room temperature for 15 min, when ethaneboronic acid (1.0g) and an aqueous solution of sodium carbonate (1.0M, 8.5 mL) are added sequentially. The mixture is heated to 75°C with stirring for 12 h, then diluted with 0.1N sodium hydroxide
30 and extracted twice with 1:1 hexane-ethyl ether. Combined extracts are dried (sodium

sulfate), filtered, concentrated, and chromatographed on silica (10:1 hexane-EtOAc) to give the title compound as a yellow oil (460 mg, 55%): ^1H NMR (CDCl_3 , 400 MHz) δ 0.95 (t, 6H), 1.28(t, 3H), 1.54(m, 2H), 1.67(m, 2H), 2.20(s, 3H), 2.65(q, 2H), 4.13(m, 2H), 7.27(d, 1H), 7.31(d, 1H), 7.48(s, 1H).LC-MS: 352 (M+1).

5

Example 84a**5-(2,4-dichlorophenyl)-3-ethoxy-6-ethyl-[N-(1-ethyl)propyl]pyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; $\text{R}_1=\text{OCH}_2\text{CH}_3$; $\text{R}_2=-\text{NHCH}(\text{CH}_2\text{CH}_3)_2$; $\text{R}_3=\text{CH}_2\text{CH}_3$]

The same reaction as in Example 83 with sodium ethoxide gives the title compound: ^1H NMR (CDCl_3 , 400 MHz) δ 0.95 (t, 6H), 1.10 (t, 3H), 1.37 (t, 3H), 1.55 (m, 2H), 1.68 (m, 2H), 2.36 (m, 2H), 4.05 (m, 1H), 4.33 (q, 2H), 4.81 (d, 1H), 7.24 (d, 1H), 7.26 (dd, 1H), 7.46 (d, 1H); MS(CI) 382.

Example 85**5-(2,4-dichlorophenyl)-6-ethyl-[N-(1-ethyl)propyl]-3-methylpyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; $\text{R}_1=\text{CH}_3$; $\text{R}_2=-\text{NHCH}(\text{CH}_2\text{CH}_3)_2$; $\text{R}_3=\text{CH}_2\text{CH}_3$]

3-Bromo-5-(2,4-dichlorophenyl)-6-ethyl-[N-(1-ethyl)propyl] pyrazine-2-amine obtained from Example 79 is converted to the title compound by using methylmagnesium bromide according to the same procedure used in Example 21: ^1H NMR (CDCl_3 , 400 MHz) δ 0.97 (t, 6H), 1.12 (t, 3H), 1.56 (m, 2H), 1.65 (m, 2H), 2.35 (s, 3H), 2.42 (m, 2H), 4.04 (m, 2H), 7.25 (d, 1H), 7.29 (dd, 1H), 7.46 (d, 1H); MS (CI) 352.

Example 86**3-Bromo-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]-6-methoxypyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; $\text{R}_1=\text{Br}$; $\text{R}_2=-\text{NHCH}(\text{CH}_2\text{CH}_3)_2$; $\text{R}_3=\text{OCH}_3$]

6-Chloro-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]pyrazine-2-amine obtained in Example 75C is converted to 5-(2,4-dichlorophenyl)-6-methoxy-[N-(1-ethyl)propyl]pyrazine-2-amine according to the same procedure used in Example 83, which is further converted to the title compound according to the same procedure used in Example 79: ^1H NMR (CDCl_3 , 400 MHz) δ 0.97 (t, 6H), 1.60 (m, 2H), 1.70 (m, 2H), 3.89

(s, 3H), 3.92 (m, 1H), 4.98 (d, 1H), 7.27 (dd, 1H), 7.34 (d, 1H), 7.44 (d, 1H); MS (CI) 418.

Example 86a

5 5-(2,4-Dichlorophenyl)-[N-(1-ethyl)propyl]-3,6-dimethoxypyrazine-2-amine

[Formula I: Ar=2,4-dichlorophenyl; R₁=R₃=OCH₃, R₂= -NHCH(CH₂CH₃)₂]

To a solution of 3-bromo-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]-6-methoxypyrazine-2-amine (0.8 mmole, obtained in example 86) in 1-methyl-2-pyrrolidinone (5ml) was added sodium methoxide (3.0 mmole). The resulting mixture
10 was then heated to 80 °C for three days. The mixture was then diluted with water and extracted with ethyl acetate. The combined extracts were washed thoroughly with water, then brine and dried. After solvent removed, the crude was purified by silica gel column (eluted with 3% EtOAc in hexane) to give the title compound as a light yellow oil (65% yield). ¹H NMR (CDCl₃, 400 MHz) δ 0.97 (t, 6H), 1.56 (m, 2H), 1.68 (m, 2H), 3.86 (s,
15 3H), 3.94 (s,3H), 3.95(m, 1H), 4.82 (d, 1H), 7.27 (dd, 1H), 7.40 (d, 1H), 7.44 (d, 1H). MS (CI) 370.

Example 86b

3-Ethyl-5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]-6-methoxypyrazine-2-amine

20 [Formula I: Ar=2,4-dichlorophenyl; R₁=CH₂CH₃; R₃=OCH₃, R₂= -NHCH(CH₂CH₃)₂]

A: To a solution of 2-(3-pentylamino)-6-chloropyrazine (3.3g, 16.5mmol) (obtained in example 75) in 1-methyl-2-pyrrolidinone (15 mL) at room temperature is added a solution of sodium methoxide in methanol (5.0M, 10ml). The resulting solution was heated to 50°C for 20 h then evaporated and poured into water and extracted twice
25 with ethyl acetate/hexane (1:1). Combined extracts were dried (sodium sulfate), filtered, concentrated, and chromatographed to give N-(1-ethyl)propyl-6-methoxypyrazine-2-amine as a light yellow solid(98%).

B: A solution of the above solid in chloroform(60 mL) is cooled to 0°C and N-bromosuccinimide (3.0 g) is added in portions. After the addition, the mixture is further
30

stirred for 1 h while being allowed to warm to room temperature. The mixture is then diluted with dichloromethane, washed with saturated NaHCO_3 , water, brine and dried, filtered. The filtrate is concentrated and chromatographed on silica gel (dichloromethane/hexane 1:1 elution) to give 3-bromo-[N-(1-ethyl)propyl]-6-methoxypyrazine-2-amine as a yellow oil (35%). ^1H NMR (CDCl_3 , 400 MHz) δ 0.93 (t, 6H), 1.56 (m, 2H), 1.66 (m, 2H), 3.87 (s, 3H), 3.92 (m, 1H), 4.85 (d, 1H), 7.18 (s, 1H).

C: To a solution of above 3-bromo-[N-(1-ethyl)propyl]-6-methoxypyrazine-2-amine (1.27g, 4.63 mmol) in THF (30 mL) is added [1,3-bis(diphenylphosphino)propane]dichloronickel(II) (125 mg). After 10 min, ethylmagnesium bromide (1.0M in THF, 9.7 mL) is added slowly dropwise at 0 °C. The mixture is stirred at room temperature for 1 hr. The resulting dark solution is poured into aqueous ammonium chloride and extracted twice with ether. Combined extracts are dried (sodium sulfate), filtered, concentrated, and chromatographed (2.5% MeOH/ CH_2Cl_2) to give the desired product 3-ethyl-[N-(1-ethyl)propyl]-6-methoxypyrazine-2-amine as a light yellow oil (55%).


D: A solution of the above oil (0.55g, 2.46 mmol) in chloroform (10 mL) is cooled to 0°C and N-bromosuccinimide (445 mg) is added in portions. After the addition, the mixture is further stirred for 30 min while being allowed to warm to room temperature. The mixture is then concentrated to dryness in vacuo, and chromatographed on silica gel (5% ethyl acetate in hexane elution) to give 5-bromo-3-ethyl-6-methoxy-[N-(1-ethyl)propyl]pyrazine-2-amine (85%).

E. A mixed solution of the above bromide (100mg; 0.33 mmol) and tetrakis(triphenylphosphine)palladium(0) (5 mol%) in ethyleneglycol dimethyl ether (3 mL) is stirred at room temperature for 15 min, when 2,4-dichlorobenzeneboronic acid (95mg) and an aqueous solution of sodium carbonate (1.0M, 0.75 mL) are added sequentially. The mixture is heated to 75°C with stirring for 15 h, then diluted with water and extracted twice with 1:1 hexane-ethyl ether. Combined extracts are dried (sodium


sulfate), filtered, concentrated, and chromatographed on silica (6% ethyl acetate in hexane) to give the title compound as a yellow oil (121 mg, 99%): ¹H NMR (CDCl₃, 400 MHz) δ 0.97 (t,6H), 1.27(t,3H), 1.56(m,2H), 1.70(m,2H), 2.62(q,2H), 3.85(s,3H), 4.00(m, 1H), 4.18 (d,1H), 7.28(d,1H), 7.38(d,1H), 7.44(s,1H). MS: 368.

Examples 86C-86D in Table VI may be prepared following the general procedure described in Example 86b or Example 86e using the corresponding phenylboronic acids

Table VI



EX#	R ₂	R ₄	¹ H NMR	MS (CI)	NAME
86C	OCH ₃	OCH ₃	0.96 (t, 6H), 1.27 (t, 3H), 1.56 (m, 2H), 1.70 (m, 2H), 2.62 (q, 2H), 3.78 (s, 3H), 3.82 (s, 3H), 3.85 (s, 3H), 4.02 (m, 2H), 6.54 (d, 1H), 6.56 (dd, 1H), 7.26 (d, 1H).	360	3-ethyl-6-methoxy-N-(1-ethylpropyl)-5-(2,4-dimethoxyphenyl)pyrazin-2-amine
86D	OCF ₃	OCH ₃	0.97 (t, 6H), 1.27 (t, 3H), 1.56 (m, 2H), 1.70 (m, 2H), 2.62 (q, 2H), 3.83 (s, 3H), 3.86 (s, 3H), 4.01 (m, 1H), 4.10 (d, 1H), 6.85 (s, 1H), 6.87 (d, 1H), 7.43 (d, 1H).	414	3-ethyl-6-methoxy-N-(1-ethylpropyl)-5-(2-difluoromethoxy-4-methoxyphenyl)pyrazin-2-amine
86E	OCH ₃	CF ₃	0.96 (t, 6H), 1.27 (t, 3H), 1.58 (m, 2H), 1.70 (m, 2H), 2.63 (q, 2H), 3.86 (s, 3H), 3.87 (s, 3H), 4.01 (m, 1H), 4.13 (m, 1H), 7.16 (s, 1H), 7.27 (d, 1H), 7.47 (d, 1H).	398	3-ethyl-6-methoxy-N-(1-ethylpropyl)-5-(2-methoxy-4-trifluoromethylphenyl)pyrazin-2-amine

Example 86f**3-Ethyl-5-(2-methoxy-4-trifluoromethoxyphenyl)-[N-(1-ethyl)propyl]-6-methoxypyrazine-2-amine**5 **Scheme V**

- A. *(1-Ethyl-propyl)-(3-ethyl-pyrazin-2-yl)-amine*. In a pressure tube, 2-chloro-3-ethyl-pyrazine (1.4 g, 10 mmol), $\text{Pd}_2(\text{dba})_3$ (229 mg, 0.25 mmol) and $\text{P}(\text{t-Bu})_3$ (100 L, 0.4 mmol) were dissolved in toluene (15 mL) and stirred at room temperature for 30 min. 1-Ethyl-propylamine (1.75 mL, 15 mmol) and KOt-Bu (1M in THF, 15 mmol, 15 mL) were added, and the dark solution was stirred at 55°C (oil bath temperature) for 90 min. The reaction mixture was allowed to reach room temperature, diluted with ethyl ether (100 mL) and washed with brine (3 x 100 mL). After drying with MgSO_4 , the solvents were removed under reduced pressure and a dark oil obtained. Flash chromatography (100% hexanes to 20% ethyl acetate in hexanes) produced the desired product (710 mg, 37%). ^1H NMR (CDCl_3 , 300 MHz): 7.84 (1H, d, $J=8.1\text{Hz}$), 7.68 (1H, d, $J=9\text{Hz}$), 4.15 (br, 1H), 4.05 (quint, 1H), 2.6 (2H, q, $J=7.4\text{Hz}$), 1.4-1.6 (m, 4H), 1.32 (t, 3H, $J=7.4\text{Hz}$), 0.95 (t, 6H, $J=7.4\text{Hz}$). MS: 194 ($\text{M}+1$, positive mode) and 192 ($\text{M}-1$, negative mode).

B. *(5-Bromo-3-ethyl-pyrazin-2-yl)-(1-ethyl-propyl)-amine*. The product from step 1 (650 mg, 3.4 mmol) was dissolved in chloroform (20 mL) and treated at room

temperature with N-bromosuccinimide (600 mg, 3.5 mmol). After 5 min the mixture was diluted with chloroform (100 mL) and the organic solution washed with brine (3 x 100 mL), dried (MgSO₄), filtered, and the solvent removed under reduced pressure to produce 5-bromo-3-ethyl-pyrazin-2-yl)-(1-ethyl-propyl)-amine (700 mg, 77%). H-1 NMR (CDCl₃, 400 MHz): 7.94 (1H, s), 4.1 (br, 1H), 3.95 (quint, 1H), 2.60 (2H, q, J=7.4Hz), 1.4-1.6 (m, 4H), 1.32 (t, 3H, J=7.4Hz), 0.95 (t, 6H, J=7.4Hz). MS: 274.1 (M+1, positive mode) and 270.3 (M-1, negative mode).

C. *[3-Ethyl-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazin-2-yl]-(1-ethyl-propyl)-amine*. In a pressure tube, a mixture of the product from step 2 (700 mg, 2.6 mmol), 2-methoxy-4-trifluoromethoxyboronic acid (1.0 g, 4.2 mmol) and Pd(PPh₃)₄ (100 mg) in toluene (10 mL), ethanol (0.5 mL) and aqueous K₂CO₃ (2M, 5 mL) was heated in oil bath at 80°C for 16 h. The mixture was diluted with ethyl acetate, and the organic fraction washed with NaOH (2M, 50 mL) and brine (3 x 50 mL), dried (MgSO₄) filtered, and the solvent removed under reduced pressure. Flash chromatography of the residue (100% hexanes to 4% ethyl acetate in hexanes) provided the desired product as an oil (850 mg, 85%). H-1 NMR (CDCl₃, 300 MHz): 8.53 (1H, s), 7.91 (1H, d, J=8.8Hz), 6.93 (1H, d, J=8.8 Hz), 6.80 (s, 1H), 4.18 (1H, d, J=8.2Hz), 4.10 (quint, 1H, J=5.8Hz), 3.88 (3H, s), 2.6 (2H, q, J=7.4Hz), 1.4-1.6 (m, 4H), 1.37 (t, 3H, J=7.4Hz), 0.95 (t, 6H, J=7.4Hz). MS: 384.3 (M+1, positive mode) and 382.2 (M-1, negative mode).

D. *[3-Ethyl-5-(2-methoxy-4-trifluoromethoxy-phenyl)-4-oxy-pyrazin-2-yl]-(1-ethyl-propyl)-amine*. The aminopyrazine obtained in step 3 (370 mg, 0.97 mmol) was dissolved in dichloromethane (15 mL) and treated with solid *m*-chloroperoxybenzoic acid (374 mg, 1.3 eq) at room temperature. After 3 h the reaction was worked up by diluting with dichloromethane (50 mL) and washing with NaOH 2M (25 mL) and brine (3 x 50 mL). The organic solution was dried (MgSO₄), filtered and the solvent evaporated under reduced pressure. The residue was purified by preparative thin layer chromatography, eluting with 30% ethyl acetate in hexanes), furnishing the desired product (55 mg, 14%).

NMR (CDCl₃, 400 MHz) H-1: 7.86 (1H, s), 7.38 (1H, d, J=8.4Hz), 6.89 (1H, d, J=7.2 Hz), 6.81 (s, 1H), 4.21 (1H, d, J=8.0Hz), 4.08 (quint, 1H, J=6.0Hz), 3.81 (3H, s), 2.9 (2H, q, J=7.6Hz), 1.5-1.75 (m, 4H), 1.23 (t, 3H, J=7.2Hz), 0.96 (t, 6H, J=7.6Hz). C-13: 158.96, 154.43, 150.64, 142.56, 132.86, 132.47, 131.32, 119.15, 112.22, 104.58, 104.58, 56.01, 53.35, 26.96, 17.89, 10.04, 8.86. F-19 NMR: -58.04 (s). (MS: 400.3 (M+1, positive mode) and 398.3 (M-1, negative mode).

E. *[6-Chloro-3-ethyl-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazin-2-yl]-(1-ethyl-propyl)-amine*. The *N*-oxide from step 4 (40 mg, 0.1 mmol) was dissolved in OPCl₃ (1.5 mL) and heated at 80°C for 16 h. After cooling down to room temperature, the reaction mixture was diluted with ethyl ether (100 mL) and washed with NaOH (2M, 50 mL) and brine (3 x 50 mL), dried (MgSO₄), filtered, and the solvent evaporated under reduced pressure to afford the desired chloropyrazine (40 mg, 96%). NMR (CDCl₃, 400 MHz) H-1: 7.32 (1H, d, J=8.4Hz), 6.89 (1H, d, J=7.2 Hz), 6.80 (s, 1H), 4.26 (1H, d, J=8.4Hz), 4.07 (quint, 1H, J=5.6Hz), 3.82 (3H, s), 2.64 (2H, q, J=7.6Hz), 1.5-1.75 (m, 4H), 1.29 (t, 3H, J=7.6Hz), 0.96 (t, 6H, J=7.2Hz). C-13: 158.20, 150.91, 150.14, 143.80, 140.95, 134.30, 131.94, 126.07, 112.44, 104.58, 55.78, 53.01, 26.81, 25.75, 10.80, 10.02. F-19 NMR: -58.03 (s). (MS: 418.2 (M+1, positive mode) and 416.2 (M-1, negative mode).

20

F. *[3-Ethyl-6-methoxy-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazin-2-yl]-(1-ethyl-propyl)-amine*. In a pressure tube, the chloropyrazine from step 5 (30 mg) was dissolved in DMF (2 mL) and treated with sodium methoxide (100 mg) at 80°C for 120 h. The reaction mixture was diluted with ethyl ether (50 mL) and washed with brine. The residue was purified by preparative thin layer chromatography (15% ethyl acetate in hexanes to produce [3-ethyl-6-methoxy-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazin-2-yl]-(1-ethyl-propyl)-amine (10 mg, 33%), identical by tlc and NMR with an authentic specimen.

30

Example 87**6-chloro-[N-(1-ethyl)propyl]-3-methoxy-5-(2,4-dichlorophenyl)pyrazine-2-amine**

[Formula I: Ar=2,4-dichlorophenyl; R₁=OCH₃; R₂= -NHCH(CH₂CH₃)₂; R₃=Cl]

A. A solution in DMF (2 mL) of 3-bromo-6-chloro-[N-(1-ethyl)propyl]pyrazine-2-amine (0.50 g; 1.8 mmol) obtained as a minor product in Example 75B is added into a solution of sodium methoxide (freshly prepared from 90 mg of sodium metal in methanol; 3.9 mmol) in DMF (3 mL). The mixture is stirred at room temperature overnight, then poured into aqueous ammonium chloride solution and extracted twice with hexane-ethyl ether. Combined extracts are dried (sodium sulfate), filtered, concentrated *in vacuo*, and chromatographed on silica gel to give 6-chloro-3-methoxy-[N-(1-ethyl)propyl]pyrazine-2-amine (410 mg).

B. The above material is brominated according to the same procedure used in Example 79 and further converted to the title compound according to the same Suzuki coupling procedure used in Example 1C: ¹H NMR (CDCl₃, 400 MHz) δ 0.95 (t, 6H), 1.5 - 1.7 (m, 4H), 2.15 (s, 3H), 3.8 (s, 3H), 3.85 (s, 3H), 3.95 (s, 3H), 4.0 (m, 1H), 4.8 (br d, 1H), 6.55 (s, 1H), 6.55 (d, 1H), 7.3 (d, 1H).

Example 88

5-(2,4-dichlorophenyl)-[N-(1-ethyl)propyl]-3-methoxy-6-methylpyrazine-2-amine
[Formula I: Ar=2,4-dichlorophenyl; R₁=OCH₃; R₂= -NHCH(CH₂CH₃)₂; R₃=CH₃]

A. 6-chloro-3-methoxy-[N-(1-ethyl)propyl]pyrazine-2-amine obtained in Example 87A is converted to 6-methyl derivative according to the same procedure used in Example 85 and further brominated according to the same procedure used in Example 79.

B. The 5-bromopyrazine obtained as above is converted to the title compound according to the same procedure used in Example 1C: ¹H NMR (CDCl₃, 400 MHz) δ 0.95 (t, 6H), 1.5 - 1.7 (m, 4H), 2.15 (s, 3H), 3.9 (s, 3H), 4.05 (m, 1H), 4.8 (br d, 1H), 7.3 (s, 2H), 7.45 (s, 1H); MS (CI) 356.

Examples 89-90 may be prepared according to the general procedure described in Example 88.

5 **Example 89**

[N-(1-ethyl)propyl]-3-methoxy-5-(2,4-dimethoxyphenyl)-6-methylpyrazine-2-amine

[Formula I: Ar=2,4-dimethoxyphenyl; R₁=OCH₃; R₂= -NHCH(CH₂CH₃)₂; R₃=CH₃]

¹H NMR (CDCl₃, 400 MHz): δ 0.95 (t, 6H), 1.5 - 1.7 (m, 4H), 2.15 (s, 3H), 3.8 (s, 3H),
3.85 (s, 3H), 3.95 (s, 3H), 4.0 (m, 1H), 4.7 (br d, 1H), 6.5 (s, 1H), 6.55 (d, 1H), 7.2 (d,
10 1H); MS (CI) 346.

Example 90


[N-(1-ethyl)propyl]-3-methoxy-5-(4-methoxy-2-methylphenyl)-6-methylpyrazine-2-amine [Formula I: Ar=2-methyl-4-methoxyphenyl; R₁=OCH₃; R₂= -NHCH(CH₂CH₃)₂;

15 R₃=CH₃]

¹H NMR (CDCl₃, 400 MHz): δ 0.95 (t, 6H), 1.5 - 1.7 (m, 4H), 2.15 (s, 3H), 2.2 (s, 3H), 3.8 (s, 3H), 3.9 (s, 3H), 4.0 (m, 1H), 4.7 (br d, 1H), 6.75 (d, 1H), 6.8 (s, 1H), 7.15 (d, 1H); MS (CI) 330.

20 **Example 91**

{4-[6-Ethyl-5-(ethylpropoxy)-3-methoxypyrazin-2-yl]-3-methoxyphenoxy}trifluoromethane



A. To a stirred solution of 2,6-dichloropyrazine (25g, 0.167 mol) and ethanol (120 mL) in a sealed tube is added 4-methoxybenzylamine (68.6g, 0.5 mol). The mixture is heated to 115 °C for 12 h and cooled. The white solid is removed by filtration and the filtrate evaporated. The residue is dissolved in ethyl acetate and washed successively with 2M sodium hydroxide, water and aqueous sodium chloride. The organics are dried (magnesium sulfate), filtered, and evaporated to give (6-chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine (35.5 g), a white solid.

B. To a stirred solution of (6-chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine (5.0 g, 20.0 mmol) in DMF (30 mL) is added sodium methoxide (7.50 g, 125.0 mmol). The mixture is heated at reflux for 12 h, cooled and partitioned between ethyl acetate (100 mL) and water (100 mL). The layers are separated and the organic layer washed with water (3 x 100 mL). The combined extracts are dried

(magnesium sulfate), filtered, and evaporated to give [(4-methoxyphenyl)methyl](6-methoxypyrazin-2-yl)amine (4.65 g), a yellow solid.

5 C. A solution of [(4-methoxyphenyl)methyl](6-methoxypyrazin-2-yl)amine (2.45g, 10.0 mmol) in chloroform (50 mL) is cooled to 0 C and N-bromosuccinimide (1.8g, 10.0 mmol) is added in portions. After the addition, the mixture is further stirred for 1 h while being allowed to warm to room temperature. The mixture is washed with saturated aqueous sodium bicarbonate, aqueous sodium chloride, dried (magnesium sulfate), filtered and evaporated. The residue is purified by flash chromatography, eluting with
10 20% ether in hexanes to give (5-bromo-6-methoxypyrazin-2-yl)[(4-methoxyphenyl)methyl]amine (1.1 g).

D. To a stirred solution of (5-bromo-6-methoxypyrazin-2-yl)[(4-methoxyphenyl)methyl]amine (0.80 g, 2.50 mmol) and 2-methoxy-4-
15 trifluoromethoxybenzeneboronic acid (1.75 g, 7.5 mmol) in toluene (25 mL) is added tetrakis(triphenylphosphine)palladium(0) (100 mg) and potassium carbonate (2.0 M, 2.0 mL). The mixture is heated to 85 C for 8 h, cooled to room temperature, diluted with 2.0 M sodium hydroxide and extracted twice with 1:1 hexane-ethyl ether. The combined
20 extracts are dried (sodium sulfate), filtered and concentrated. The residue is purified by flash chromatography, eluting with 60% hexanes in ether to give {6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-yl}[(4-methoxyphenyl)methyl]amine (967 mg).

E. A solution of {6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-
25 2-yl}[(4-methoxyphenyl)methyl]amine (870 mg, 2.0 mmol) in chloroform (10 mL) is cooled to 0 C and N-bromosuccinimide (356 mg, 2.0 mmol) is added in portions. After the addition, the mixture is further stirred for 1 h while being allowed to warm to room temperature. The mixture is washed with saturated aqueous sodium bicarbonate, aqueous sodium chloride, dried (magnesium sulfate), filtered, and evaporated to give {3-

bromo-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-yl}[(4-methoxyphenyl)methyl]amine (920 mg), a yellow solid.

5 F. To a stirred solution of {3-bromo-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-yl}[(4-methoxyphenyl)methyl]amine (514 mg, 1.0 mmol) and ethylboronic acid (219 mg, 3.0 mmol) in toluene (8 mL) is added tetrakis(triphenylphosphine)palladium(0) (50 mg) and potassium carbonate (2.0 M, 1.0 mL). The mixture is heated to 85 C for 8 h, cooled to room temperature, diluted with 2.0 M sodium hydroxide and extracted twice with 1:1 hexane- ethyl ether. The combined
10 extracts are dried (sodium sulfate), filtered and concentrated. The residue is purified by flash chromatography, eluting with 20% ether in hexanes to give {3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-yl}[(4-methoxyphenyl)methyl]amine (430 mg).


15 G. To {3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-yl}[(4-methoxyphenyl)methyl]amine (115 mg, 0.25 mmol) in methanol (3 mL) under a nitrogen atmosphere is added 1M hydrochloric acid in ether (2 mL) and 10% palladium on carbon (40 mg). The mixture is then hydrogenated at 1 ATM for 18 h, filtered through Celite and evaporated to give 3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-ylamine (80 mg).
20

H. To a stirred solution of 3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-ylamine (86 mg, 0.25 mmol) in 48% hydrogen bromide (0.3 mL) at 0 C is added a solution of sodium nitrite (21 mg, 0.3 mmol) in water
25 (1 mL). After 1.5 h copper bromide (43 mg, 0.3 mmol) is added and the mixture heated to 70 C for 1 h. The mixture is cooled to room temperature and extracted with ether. The extracts are dried (sodium sulfate), filtered and concentrated to give [4-(5-bromo-6-ethyl-3-methoxypyrazin-2-yl)-3-methoxyphenoxy]trifluoromethane (76 mg).

I. To a stirred solution of 3-pentanol (88 mg, 1 mmol) in THF (1 mL) is added 60% sodium hydride (12 mg, 0.3 mmol) and after 0.5 h [4-(5-bromo-6-ethyl-3-methoxypyrazin-2-yl)-3-methoxyphenoxy]trifluoromethane (41 mg, 0.1 mmol) is added. The mixture is heated to 50 C for 12 h, cooled and partitioned between ethyl acetate and
5 water. The organic layer is washed with water, brine, dried (sodium sulfate), filtered and concentrated. The residue is purified by preparative TLC eluting with 50% ether in hexanes to give {4-[6-Ethyl-5-(ethylpropoxy)-3-methoxypyrazin-2-yl]-3-methoxyphenoxy}trifluoromethane, a colorless oil (19 mg). NMR (CDCL₃, 400 MHz)
10 0.98 (t, 6H), 1.22 (t, 3H), 1.78 (m, 4H) 2.80 (q, 2H), 3.80 (s, 3H), 3.88 (s, 3H), 5.05 (quintet, 1H), 6.8 (s, 1H), 6.90 (d, 1H), 7.37 (d, 1H); MS 345 (M+1).

Additional compounds prepared and assayed by the methods described herein are shown in Table VII.

Table VII




Formula I

Cpnd No.	Ar	R ₁	R ₂	R ₃	Synthetic Method	IMR32 Binding	Name
91a	2,4,6-trichlorophenyl	CH ₃	3-pentyl amino	CH ₃	Example I	40 nM	3,6-dimethyl-N-(1-ethylpropyl)-5-(2,4,6-trimethylphenyl)pyrazin-2-amine
91b	2,4-dichlorophenyl	Br	3-pentyl amino	Cl	Example I	28 nM	3-bromo-6-chloro-N-(1-ethylpropyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine

Example 92**Synthesis of [3-ethyl-5-(2-methoxy-4-trifluoromethoxy-phenyl)-6-methylsulfanyl-pyrazin-2-yl]-(1-ethyl-propyl)-amine.**

In a pressure tube equipped with Teflon O-ring were combined [6-chloro-3-ethyl-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazin-2-yl]-(1-ethyl-propyl)-amine (100 mg) obtained in step E of Example 86e, NaSMe (200 mg), THF (5 mL) and DMF (3 mL). The mixture was heated at 80 °C (oil bath temperature) for 72 h. The crude mixture was diluted with ethyl acetate (40 mL) and water (40 mL), and the organic phase washed with brine (3 x 100 mL). After drying (MgSO₄), filtration and elimination of solvents at reduced pressure, the title compound was isolated as a clear oil by preparative thin layer chromatography (10% EtOAc in hexanes). Yield was 50 mg (49%). NMR (CDCl₃, 400 MHz) H-1: 7.32 (1H, d, J=8.4Hz), 6.88 (1H, m), 6.78 (s, 1H), 4.18 (1H, d), 4.07 (m, 1H), 3.78 (3H, s), 2.64 (2H, q), 2.24 (s, 3H), 1.5-1.75 (m, 4H), 1.25 (t, 3H), 0.96 (t, 6H). MS: 430.2 (M+1, positive mode) and 428.4 (M-1, negative mode).

Example 93**2-sec-butylsulfanyl-5-(2,4-dimethoxy-phenyl)-3,6-diethyl-pyrazine**

NaH (60 mg, 1.5 mmol, 60% in mineral oil) was added to a solution of butane-2-thiol (170 µL, 1.5 mmol) in THF (5 mL). After 10 minutes, 2-chloro-5-(2,4-dimethoxy-

phenyl)-3,6-diethyl-pyrazine (100 mg, 0.33 mmol) obtained as in Example 45, steps A-C, in THF (1 mL) was added dropwise, and the mixture heated at 80 °C (oil bath temperature) for 16 h. After extractive work-up, preparative thin layer chromatography (hexanes) furnished the title compound as a clear oil (60 mg, 51%). NMR (CDCl₃, 400 MHz) H-1: 7.19 (1H, d, J=8.4Hz), 6.58 (1H, dd, J=2.4, 8.4Hz), 6.60 (s, 1H, J=2.4Hz), 3.99 (sext, 1H, J=6.8Hz), 3.85 (3H, s), 3.75 (3H, s), 2.81 (2H, q, J=7.4Hz), 2.58 (2H, br q, J=6.8Hz), 1.6-1.9 (m, 4H), 1.44 (3H, d, J=6.4Hz), 1.28 (t, 3H, J=7.6Hz), 1.19 (3H, t, J=7.6Hz), 1.06 (t, 3H, J=7.2Hz). C-13: 161.10, 157.86, 153.49, 151.79, 151.67, 144.47, 131.68, 121.18, 104.78, 98.60, 55.44, 55.33, 40.87, 29.65, 27.54, 27.08, 20.56, 12.43, 12.09, 11.51. MS: 414.2 (M+1, positive mode) and 412.2 (M-1, negative mode).

Example 93a

2-Sec-butylsulfanyl-5-(2,4-dichloro-phenyl)-3,6-diethyl-pyrazine.

By a similar procedure, but starting from 2-chloro-5-(2,4-dichloro-phenyl)-3,6-diethyl-pyrazine obtained as in Example 49 steps A-C, was obtained 2-sec-butylsulfanyl-5-(2,4-dichloro-phenyl)-3,6-diethyl-pyrazine. NMR (CDCl₃, 400 MHz) H-1: 7.19 (1H, d, J=2.0Hz), 7.26 (1H, dd, J=2.0, 8.0Hz), 7.18 (1H, d, J=8.0Hz), 3.92 (1H, sext, J=6.8Hz), 2.72 (2H, q, J=7.6Hz), 2.58 (2H, br), 1.6-1.8 (m, 4H), 1.36 (3H, d, J=6.8Hz), 1.20 (t, 3H, J=7.6Hz), 1.12 (3H, t, J=7.6Hz), 0.98 (t, 3H, J=7.6Hz). C-13: 153.93, 152.40, 152.00, 143.69, 136.46, 134.73, 134.40, 131.94, 129.48, 127.24, 41.01, 29.56, 27.40, 27.10, 20.45, 12.48, 11.86, 11.51. MS: 369.2 (M+1, positive mode).


Example 93b

2-Sec-butylsulfanyl-3-ethyl-6-methoxy-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazine.

By a similar procedure, but starting from 2-bromo-3-ethyl-6-methoxy-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazine obtained as in Example 49 steps A-C, was obtained 2-sec-butylsulfanyl-3-ethyl-6-methoxy-5-(2-methoxy-4-trifluoromethoxy-phenyl)-pyrazine. NMR (CDCl₃, 400 MHz) H-1: 7.37 (1H, d, J=8.4Hz), 6.91 (1H, m), 6.80 (1H, s), 3.94 (3H, s), 3.89 (1H, m), 3.79 (3H, s), 2.78 (2H, q, J=7.6Hz), 1.7-1.9 (m,


4H), 1.46 (3H, d, J=6.8Hz), 1.27 (t, 3H, J=7.2Hz), 1.07 (3H, t, J=7.2Hz). C-13: 158.43, 155.85, 150.32, 149.90, 146.33, 135.34, 133.73, 131.85, 124.81, 112.66, 104.85, 55.93, 53.57, 41.47, 29.61, 26.83, 20.61, 12.30, 11.54. F-19 NMR: -58.04 (s). MS: 417.2 (M+1, positive mode).

5

Example 94**1-[3,6-diethyl-5-(2-methylbutyl)pyrazin-2-yl]-2,4-dimethoxybenzene**

To a solution of 2-methyl-1-butene (210 mg, 3.0 mmol) in THF is added a
 10 solution of 9-BBN (9-borabicyclo[3.3.1]nonane) in THF (0.5 M, 6.0 mL, 3.0 mmol). The mixture is heated at reflux, under a nitrogen atmosphere for 12 h and cooled. To the solution is added 1-(5-bromo-3,6-diethylpyrazin-2-yl)-2,4-dimethoxybenzene (664 mg, 2.0 mmol), obtained as in Example 49 steps A-C, tetrakis(triphenylphosphine) palladium(0) (50 mg) and sodium hydroxide (3.0 M, 3.0 mL, 3.0 mmol). The mixture is
 15 heated at 50 °C for 12 h and cooled. 30% Hydrogen peroxide (1 mL) is added, the solution stirred for 1 h and the reaction mixture extracted with ether. The combined extracts are dried (sodium sulfate), filtered and concentrated. The residue is purified by flash chromatography, eluting with 40% ether in hexanes to give 1-[3,6-diethyl-5-(2-methylbutyl)pyrazin-2-yl]-2,4-dimethoxybenzene (393 mg).

20 Example 95**2-(2-methoxy-5-trifluoromethoxyphenyl)-3-ethyl-6-methylamino-5-(1-ethylpropoxy)-pyrazine**



(6-Chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine is prepared by Step A of Example 91.

5

A. A solution of 6-(Chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine in chloroform (4 mL/mmol NBS) is cooled to 0°C and N-bromosuccinimide (1.05 eq) is added in portions while stirring the reaction mixture. After complete addition, the mixture is further stirred for 1 hour while being allowed to warm to room temperature.

10 The mixture is then diluted with dichloromethane, washed with saturated NaHCO₃, water, brine and then dried and filtered. The filtrate is concentrated and purified by chromatography on silica gel to give (5-bromo-6-chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine.

15

B. In a pressure tube, a mixture of the product from step A (1 equivalent), 2-methoxy-5-trifluoromethoxyphenylboronic acid (1.6 equivalents) and Pd(PPh₃)₄ (0.04

equivalents) in toluene (4 ml/mmol of product from step A), ethanol (.2 ml/ mmol of product from step A) and aqueous K_2CO_3 (2M, 2 ml/ mmol of product from step A) is heated to 80°C for 16 h. The mixture was diluted with ethyl acetate, and the organic fraction washed with NaOH (2M, 20 mL/ mmol of product from step A) and brine (3 x 20 ml/ mmol of product from step A), then dried ($MgSO_4$), filtered and the solvent removed under reduced pressure. Chromatography of the residue provided (3-(2-methoxy-5-trifluoromethoxyphenyl)-6-chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine.

C. (5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazin-2-yl)[(4-methoxyphenyl)methyl]amine can be prepared by the method of Example 1, step A by substituting (5-(2-methoxy-5-trifluoromethoxyphenyl)-6-chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine for 2-chloro-3,6-dimethylpyrazine.

D. (5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazin-2-yl)amine can be prepared by the hydrogenation method of Example 91, step G by substituting (5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazin-2-yl)[(4-methoxyphenyl)methyl]amine for {3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-yl}[(4-methoxyphenyl)methyl]amine.

E. 2-Bromo-5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazine can be prepared by the halogenation method of Example 91, step H by substituting (5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazin-2-yl)amine for {3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-yl} amine.

F. 2-(3-Pentoxy)-(5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazine can be prepared by the method of Example 91, step I by substituting 2-bromo-(5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazin-2-yl)[(4-methoxyphenyl)methyl]amine for [4-(5-bromo-6-ethyl-3-methoxypyrazin-2-yl)-3-methoxyphenoxy]trifluoromethane.

G. 2-(3-Pentoxy)-3-bromo-(5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazine can be prepared by the method of step A of the present Example by substituting 2-(3-Pentoxy)-3-bromo-(5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazine for (6-chloropyrazin-2-yl)[(4-methoxyphenyl)methyl]amine.

5

H. To a mixture of 2-(3-Pentoxy)-3-bromo-(5-(2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazine (1 equivalent), tris(dibenzylideneacetone)dipalladium(0) (2 mol%), and 1,1'-binaphthyl-2,2'-di(diphenylphosphine) (BINAP) (6 mol%) in ethyleneglycol dimethyl ether (2.4 mL/mmol substrate) under nitrogen is added methylamine (1.2 equivalents) followed by sodium *tert*-butoxide (1.5 equivalents). The mixture is stirred at 70-80°C for about 2.5 hours, diluted with aqueous ammonium chloride, and extracted with 1:1 hexane-diethyl ether. The combined extracts are dried (sodium sulfate), filtered, concentrated and then purified by chromatography on silica gel to afford 2-(3-Pentoxy)-3-(N-methylamino)-(5-
10 (2-methoxy-5-trifluoromethoxyphenyl)-6-ethylpyrazin
15

Example 96

Assay for CRF Receptor Binding Activity

As discussed above, the following assay is defined herein as a standard in vitro
20 CRF receptor binding assay.

The pharmaceutical utility of compounds of this invention is indicated by the following assay for CRF1 receptor activity. The CRF receptor binding is performed using a modified version of the assay described by Grigoriadis and De Souza (*Methods in Neurosciences*, Vol. 5, 1991). IMR-32 human neuroblastoma cells, a cell-line that
25 naturally expresses the CRF1 receptor, are grown to confluency in DMEM containing FBS.

To prepare receptor containing membranes cells are homogenized in wash buffer (50 mM Tris HCl, 10 mM MgCl₂, 2 mM EGTA, pH 7.4) and centrifuged at 48,000 x g for 10 minutes at 4°C. The pellet is re-suspended in wash buffer and the homogenization
30 and centrifugation steps are performed two additional times.

Membrane pellets containing CRF receptors are re-suspended in 50 mM Tris buffer pH 7.7 containing 10 mM MgCl₂ and 2 mM EDTA and centrifuged for 10 minutes at 48000g. Membranes are washed again and brought to a final concentration of 1500 mg/ml in binding buffer (Tris buffer above with 0.1 % BSA, 15 mM bacitracin and 0.01 mg/ml aprotinin.). For the binding assay, 100 ml of the membrane preparation are added to 96 well microtube plates containing 100 ml of ¹²⁵I-CRF (SA 2200 Ci/mmol, final concentration of 100 pM) and 50 ml of test compound. Binding is carried out at room temperature for 2 hours. Plates are then harvested on a Brandel 96 well cell harvester and filters are counted for gamma emissions on a Wallac 1205 Betaplate liquid scintillation counter. Non specific binding is defined by 1 mM cold CRF. IC₅₀ values are calculated with the non-linear curve fitting program RS/1 (BBN Software Products Corp., Cambridge, MA). The binding affinity for the compounds of Formula I expressed as IC₅₀ value, generally ranges from about 0.5 nanomolar to about 10 micromolar. Preferred compounds of Formula I exhibit IC₅₀ values of less than or equal to 1.5 micromolar, more preferred compounds of Formula I exhibit IC₅₀ values of less than 500 nanomolar, still more preferred compounds of Formula I exhibit IC₅₀ values of less than 100 nanomolar, and most preferred compound of Formula I exhibit IC₅₀ values of less than 10 nanomolar. The compounds shown in Examples 1-95 and Examples 100-396 have been tested in this assay and found to exhibit IC₅₀ values of less than or equal to 1.5 micromolar.

Example 97

Preparation of radiolabeled probe compounds of the invention

The compounds of the invention are prepared as radiolabeled probes by carrying out their synthesis using precursors comprising at least one atom that is a radioisotope. The radioisotope is preferably selected from of at least one of carbon (preferably ¹⁴C), hydrogen (preferably ³H), sulfur (preferably ³⁵S), or iodine (preferably ¹²⁵I). Such radiolabeled probes are conveniently synthesized by a radioisotope supplier specializing in custom synthesis of radiolabeled probe compounds. Such suppliers include Amersham Corporation, Arlington Heights, IL; Cambridge Isotope Laboratories, Inc. Andover, MA; SRI International, Menlo Park, CA; Wizard Laboratories, West

Sacramento, CA; ChemSyn Laboratories, Lexena, KS; American Radiolabeled Chemicals, Inc., St. Louis, MO; and Moravek Biochemicals Inc., Brea, CA.

5 Tritium labeled probe compounds are also conveniently prepared catalytically via platinum-catalyzed exchange in tritiated acetic acid, acid-catalyzed exchange in tritiated trifluoroacetic acid, or heterogeneous-catalyzed exchange with tritium gas. Such preparations are also conveniently carried out as a custom radiolabeling by any of the suppliers listed in the preceding paragraph using the compound of the invention as substrate. In addition, certain precursors may be subjected to tritium-halogen exchange
10 with tritium gas, tritium gas reduction of unsaturated bonds, or reduction using sodium borotritide, as appropriate.

Example 98

Receptor autoradiography

15 Receptor autoradiography (receptor mapping) is carried out in vitro as described by Kuhar in sections 8.1.1 to 8.1.9 of Current Protocols in Pharmacology (1998) John Wiley & Sons, New York, using radiolabeled compounds of the invention prepared as described in the preceding Example.

20 Example 99

Additional Aspects of Preferred Compounds of the Invention

The most preferred compounds of the invention are suitable for pharmaceutical use in treating human patients. Accordingly, such preferred compounds are non-toxic. They do not exhibit single or multiple dose acute or long-term toxicity, mutagenicity (e.g., as
25 determined in a bacterial reverse mutation assay such as an Ames test), teratogenicity, tumorigenicity, or the like, and rarely trigger adverse effects (side effects) when administered at therapeutically effective dosages.

Preferably, administration of such preferred compounds of the invention at certain doses (e.g., doses yielding therapeutically effective in vivo concentrations or preferably
30 doses of 10, 50, 100, 150, or 200 mg/kg -- preferably 150 mg/kg -- administered

parenterally or preferably orally) does not result in prolongation of heart QT intervals (i.e., as determined by electrocardiography, e.g., in guinea pigs, minipigs or dogs). When administered daily for 5 or preferably ten days, such doses of such preferred compounds also do not cause liver enlargement resulting in an increase of liver to body weight ratio of more than 100%, preferably not more than 75% and more preferably not more than 50% over matched controls in laboratory rodents (e.g., mice or rats). In another aspect such doses of such preferred compounds also preferably do not cause liver enlargement resulting in an increase of liver to body weight ratio of more than 50%, preferably preferably not more than 25%, and more preferably not more than 10% over matched untreated controls in dogs or other non-rodent animals.

In yet another aspect such doses of such preferred compounds also preferably do not promote the release of liver enzymes (e.g., ALT, LDH, or AST) from hepatocytes in vivo. Preferably such doses do not elevate such enzymes by more than 100%, preferably not by more than 75% and more preferably not by more than 50% over matched untreated controls in laboratory rodents. Similarly, concentrations (in culture media or other such solutions that are contacted and incubated with cells in vitro) equivalent to two, fold, preferably five-fold, and most preferably ten-fold the minimum in vivo therapeutic concentration do not cause release of any of such liver enzymes from hepatocytes in vitro.

Because side effects are often due to undesirable receptor activation or antagonism, preferred compounds of the invention exert their receptor-modulatory effects with high selectivity. This means that they do not bind to certain other receptors (i.e., other than CRF receptors) with high affinity, but rather only bind to, activate, or inhibit the activity of such other receptors with affinity constants of greater than 100 nanomolar, preferably greater than 1 micromolar, more preferably greater than 10 micromolar and most preferably greater than 100 micromolar. Such receptors preferably are selected from the group including ion channel receptors, including sodium ion channel receptors, neurotransmitter receptors such as alpha- and beta-adrenergic receptors, muscarinic receptors (particularly m1, m2, and m3 receptors), dopamine receptors, and metabotropic glutamate receptors; and also include histamine receptors and cytokine receptors, e.g.,

interleukin receptors, particularly IL-8 receptors. The group of other receptors to which preferred compounds do not bind with high affinity also includes GABA_A receptors, bioactive peptide receptors (including NPY and VIP receptors), neurokinin receptors, bradykinin receptors (e.g., BK1 receptors and BK2 receptors), and hormone receptors (including thyrotropin releasing hormone receptors and melanocyte-concentrating hormone receptors).

Example 99a

Absence of Sodium Ion Channel Activity

Preferred compounds of the invention do not exhibit activity as Sodium ion channel blockers. Sodium channel activity may be measured a standard *in vitro* sodium channel binding assays such as the assay given by Brown et al. (J. Neurosci. (1986) 265: 17995-18004). Preferred compounds of the invention exhibit less than 15 percent inhibition, and more preferably less than 10 percent inhibition, of sodium channel specific ligand binding when present at a concentration of 4 uM. The sodium ion channel specific ligand used may be labeled batrachotoxinin, tetrodotoxin, or saxitoxin. Such assays, including the assay of Brown referred to above, are performed as a commercial service by CEREP, INC., Redmond, WA.

Alternatively, sodium ion channel activity may be measured *in vivo* in an assay of anti-epileptic activity. Anti-epileptic activity of compounds may be measured by the ability of the compounds to inhibit hind limb extension in the supra maximal electro shock model. Male Han Wistar rats (150-200mg) are dosed i.p. with a suspension of 1 to 20 mg of test compound in 0.25% methylcellulose 2 hr. prior to test. A visual observation is carried out just prior to testing for the presence of ataxia. Using auricular electrodes a current of 200 mA, duration 200 millisec, is applied and the presence or absence of hind limb extension is noted. Preferred compounds of the invention do not exhibit significant anti-epileptic activity at the $p < 0.1$ level of significance or more preferably at the $p < 0.05$ level of significance as measured using a standard parametric assay of statistical significance such as a student's T test.

Example 99b**Optimal in vitro half-life**

Compound half-life values ($t_{1/2}$ values) may be determined via the following standard liver microsomal half-life assay. Liver microsomes obtained from pooled liver samples and prepared so that the P-450 enzyme content is approximately 0.5 nmol/ mg protein. Reactions are preformed in a 5ml well deep-well plate as follows:

Phosphate buffer: 19 mL 0.1 M NaH_2PO_4 , 81 mL 0.1 Na_2HPO_4 , pH 7.4 with H_3PO_4 .

CoFactor Mixture: 16.2 mg NADP, 45.4 mg Glucose-6-phosphate in 4 mL 100 mM MgCl_2 . Glucose-6-phosphate dehydrogenase: 214.3 ul glucose-6-phosphate dehydrogenase, 1285.7 ul distilled water

Starting Reaction Mixture: 3 mL CoFactor Mixture, 1.2 mL Glucose-6-phosphate dehydrogenase 6 identical samples wells each containing 25 ul microsomes, 5 ul of test compound (from a 100 uM stock), and 399 ul 0.1 M phosphate buffer, pH 7.4, are prepared. A seventh well containing 25 ul microsomes, 399 ul 0.1 M phosphate buffer, pH 7.4, and 5 ul (from a 100 uM stock) of a compound, e.g. DIAZEPAM, CLOZEPINE, with known metabolic properties is used as a positive control. Reactions are preincubated at 39 °C for 10 minutes. 71 ul Starting Reaction Mixture is added to 5 of the 6 reaction wells and to the positive control well, 71 ul 100 mM MgCl_2 is added to the sixth reaction well, which is used as a negative control. At each time point (0, 1, 3, 5, and 10 minutes) 75 ul reaction is pipetted into a 96-well deep-well plate reaction well containing 75 ul ice-cold acetonitrile. Samples are vortexed and centrifuged 10 minutes at 6000 rpm (Sorval T 6000D rotor). Supernatant, 75 ul from each reaction well, is transferred to a 96-well plate containing 150 ul internal standard per well. The remaining test compound is quantitated via LCMS and Compound concentration vs time is plotted and commercially available statistical software is used to extrapolate to the $t_{1/2}$ value of the test compound.

Preferred compounds of the invention exhibit in vitro $t_{1/2}$ values of greater than 10 minutes and less than 4 hours. Most preferred compounds of the invention exhibit in vitro $t_{1/2}$ values of between 30 minutes and 1 hour in human liver microsomes.

Example 99c**MDCK Toxicity**

Toxicity of a test compound may be assessed by measuring the effect of the compound on ATP production by MDCK cells.

5 MDCK cells, product no. CCL-34, are purchased from ATCC, Manassas, VA and maintained in sterile conditions by the methods described in the supplier's production information sheet. The PACKARD BIOSCIENCE, (Groningen, The Netherlands) ATP-LITE-M Luminescent ATP detection kit, product no. 6016941, may be used to monitor ATP production in MDCK cells.

10 Prior to assay 1 ul of test compound or control sample is pipetted into PACKARD (Meriden, CT) clear bottom 96-well plates. Test compounds and control samples are diluted in DMSO to give final concentration in the assay of 10 micromolar, 100 micromolar, or 200 micromolar. Control samples are drug compounds having known toxicity properties.

15 Confluent MDCK cells are trypsinized, harvested, and diluted to a concentration of 0.1×10^6 cells/ ml with warm ATCC Eagle's Minimum Essential Medium (catalog # 30-2003). Warm Eagle's MEM without cells (100ul) is pipetted in five wells a 96-well plate. These wells are used to determine the standard curve. Cells in Eagle's MEM (100ul or 10,000 cells) are pipetted into the remaining wells of the 96-well plates. All
20 samples are incubated at 37 °C under carbogen (95% O₂, 5% CO₂) for 2 hours with constant shaking. After incubation 50 ul mammalian cell lysis solution is added to well of the 96-well plates, wells are covered with Packard topseal stickers, and plates are shaken at approximately 700 rpm on a microtiter shaker for 2 minutes.

During the incubation, Packard ATP Lite-M reagents are allowed to equilibrate to
25 room temperature. Once equilibrated the lyophilized substrate solution is reconstituted in 5.5 mls of substrate buffer solution (from kit). Lyophilized ATP standard solution is reconstituted in deionized water to give a 10 mM stock. Standard (10 ul), diluted so that a 10 ul aliquot yields a final concentration of 200 nM, 100 nM, 50 nM, 25 nM, or 12.5 nM is added to each of the five standard curve wells, which do not contain cells.

30 Substrate solution (50 ul) is added to all wells. Wells are covered with Packard







- topseal stickers, and plates are shaken at approximately 700 rpm on a microtiter shaker for 2 minutes. A white Packard sticker is attached to the bottom of each plates and samples are dark adapted by wrapping plates in foil and placing in the dark for 10 minutes. Luminescence is then quantitated at 22 °C using a luminescence counter, e.g.
- 5 Packard TopCount Microplate Scintillation and Luminescence Counter or Tecan Spectrafluor plus.

- Luminscence values at each concentration are compared to the values computed from the standard curve for that concentration. Preferred test compounds exhibit luminescence values 80 % or more of the standard, or preferably 90 % or more of the
- 10 standard, when 10 micromolar concentration of the test compound is used. When a 100 micromolar concentraion of the test compound is used, preferred test compounds exhibit luminescence values 50 % or more of the standard, or more preferably 80 % or more of the standard.







Examples 100-396

Additional compounds of the invention prepared by the methods described for compounds shown in Examples 1-95 are shown in TABLE VIII







EX#	STRUCTURE	IUPAC Name
100		5-(2,4-dimethoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
101		6-ethyl-N-(1-ethylpropyl)-3-methoxy-5-[2-methoxy-4-(trifluoromethyl)phenyl]pyrazin-2-amine
102		3-bromo-5-(2,4-dimethoxyphenyl)-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
103		6-ethyl-N-(1-ethylpropyl)-3-methoxy-5-[4-methoxy-2-(trifluoromethoxy)phenyl]pyrazin-2-amine
104		5-[4-bromo-2-(trifluoromethoxy)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine







105		6-ethyl-N-(1-ethylpropyl)-3-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl] pyrazin-2-amine
106		2,5-diethyl-3-(1-ethylpropoxy)-6-[4-methoxy-2-(trifluoromethoxy)phenyl] pyrazine
107		5-(2,4-dimethoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)-N-propylpyrazin-2-amine
108		3-ethyl-N-(1-ethylpropyl)-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl] pyrazin-2-amine
109		5-(2,4-dimethoxyphenyl)-N-(1-ethylpropyl)-3,6-dimethoxypyrazin-2-amine
110		5-(5-chloro-4-methoxy-2-methylphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine







111		5-(5-chloro-4-methoxy-2-methylphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
112		5-(4-chloro-2-methylphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
113		5-[2-chloro-4-(trifluoromethyl)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
114		5-(5-bromo-2,4-dimethoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
115		5-(2,4-dichloro-6-methylphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
116		3,6-diethyl-N-(1-ethylpropyl)-5-(2,3,4-trimethoxyphenyl)pyrazin-2-amine







117		5-[2-chloro-4-(trifluoromethyl)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
118		5-[4-(difluoromethoxy)-2-methoxyphenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
119		2-[2-chloro-4-(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
120		2-[2,4-bis(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
121		5-(2,4-difluorophenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
122		3,6-diethyl-N-(1-ethylpropyl)-5-(1-naphthyl)pyrazin-2-amine







123		3,6-diethyl-N-(1-ethylpropyl)-5-(2-naphthyl)pyrazin-2-amine
124		6-ethyl-N-(1-ethylpropyl)-3-methoxy-5-[4-methoxy-2-(trifluoromethyl)phenyl]pyrazin-2-amine
125		5-(2,4-dimethoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)-N-methylpyrazin-2-amine
126		5-[4-chloro-2-(trifluoromethyl)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
127		5-[2-chloro-4-(trifluoromethoxy)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
128		2-(2,4-dichlorophenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine







129		2-(2,4-dichlorophenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
130		2-(2,4-dichlorophenyl)-3,6-diethyl-5-(1-ethyl-2-methylpropoxy)pyrazine
131		3,6-diethyl-N-(1-ethylpropyl)-5-[2-methoxy-6-(trifluoromethoxy)phenyl]pyrazin-2-amine
132		2-(2,4-dichlorophenyl)-3,6-diethyl-5-(1-phenylpropoxy)pyrazine
133		2-(2,4-dichlorophenyl)-3,6-diethyl-5-[(1-ethylpentyl)oxy]pyrazine
134		5-[2,6-dichloro-4-(trifluoromethyl)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine







135		5-(2-chloro-4-methoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
136		5-(2-chloro-4-methoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
137		3,6-diethyl-N-(1-ethylpropyl)-5-[4-fluoro-2-(trifluoromethyl)phenyl]pyrazin-2-amine
138		5-[2,4-bis(trifluoromethyl)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
139		5-(2,6-dichloro-4-methoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
140		5-[2-chloro-4-(trifluoromethyl)phenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine







141		2-[cyclopropyl(phenyl)methoxy]-5-(2,4-dichlorophenyl)-3,6-diethylpyrazine
142		5-[2-chloro-4-(trifluoromethoxy)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
143		3,6-diethyl-N-(1-ethylpropyl)-5-(3-methyl-1,1'-biphenyl-4-yl)pyrazin-2-amine
144		2-[2-chloro-4-(trifluoromethoxy)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
145		3,6-diethyl-N-(1-ethylpropyl)-5-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
146		6-ethyl-N-(1-ethylpropyl)-3-methoxy-5-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazin-2-amine







147		5-[2,4-bis(trifluoromethyl)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
148		5-(2-chloro-4-ethoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
149		2-(2,4-dimethoxyphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
150		2-sec-butoxy-5-(2,4-dichlorophenyl)-3,6-diethylpyrazine
151		2-(2,4-dichlorophenyl)-3,6-diethyl-5-(1-methylbutoxy)pyrazine
152		2-(2,4-dichlorophenyl)-5-(1,2-dimethylpropoxy)-3,6-diethylpyrazine







153		2-(2,4-dichlorophenyl)-3,6-diethyl-5-(1-propylbutoxy)pyrazine
154		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[4-methoxy-2-(trifluoromethoxy)phenyl]pyrazine
155		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[4-methoxy-2-(trifluoromethyl)phenyl]pyrazine
156		2-[4-chloro-2-(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
157		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazine
158		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-(4-methoxy-2-methylphenyl)pyrazine







159		N-(1-ethylpropyl)-3-methoxy-5-[4-methoxy-2-(trifluoromethoxy)phenyl]-6-methylpyrazin-2-amine
160		N-(1-ethylpropyl)-3-methoxy-5-[4-methoxy-2-(trifluoromethyl)phenyl]-6-methylpyrazin-2-amine
161		5-[4-chloro-2-(trifluoromethyl)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
162		N-(1-ethylpropyl)-3-methoxy-6-methyl-5-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
163		N-(1-ethylpropyl)-3-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]-6-methylpyrazin-2-amine
164		3,6-diethyl-N-(1-ethylpropyl)-5-[4-methyl-2-(trifluoromethyl)phenyl]pyrazin-2-amine

165		6-ethyl-N-(1-ethylpropyl)-3-methoxy-5-[4-methyl-2-(trifluoromethyl)phenyl]pyrazin-2-amine
166		5-(4-tert-butyl-2,6-dimethylphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
167		5-[4-chloro-2-(trifluoromethyl)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
168		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-(2-methoxy-4,6-dimethylphenyl)pyrazine
169		2-[2,4-bis(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
170		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[4-methyl-2-(trifluoromethyl)phenyl]pyrazine

171		3-ethyl-N-(1-ethylpropyl)-6-methoxy-5-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
172		5-(2,4-dimethylphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
173		5-(2-chloro-4-fluorophenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
174		5-[4-chloro-2-(trifluoromethyl)phenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
175		5-(2,4-dimethylphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
176		5-(2-chloro-4-fluorophenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine







177		5-(2-chloro-4-ethoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
178		N-(1-ethylpropyl)-3-methoxy-5-[2-methoxy-4-(trifluoromethyl)phenyl]-6-methylpyrazin-2-amine
179		5-(2-chloro-4-ethoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
180		5-(2,4-dimethylphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
181		3-ethyl-N-(1-ethylpropyl)-6-methoxy-5-[4-methyl-2-(trifluoromethyl)phenyl]pyrazin-2-amine
182		3-chloro-4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}phenol







183		O-(3-chloro-4-{5-[(1-isopropyl-2-methylpropoxy)-3-methoxy-6-methylpyrazin-2-yl]phenyl} S-propyl (dithiocarbonate)
184		5-[2-chloro-4-(trifluoromethoxy)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
185		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazine
186		2,5-diethyl-3-(4-fluoro-2-methoxyphenyl)-6-(1-isopropyl-2-methylpropoxy)pyrazine
187		2,5-diethyl-3-(1-ethylpropoxy)-6-[2-methoxy-4,6-bis(trifluoromethyl)phenyl]pyrazine
188		5-(2,6-dichlorophenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine

189		5-(2,6-dimethoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
190		5-(2,6-dimethylphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
191		3,6-diethyl-N-(1-ethylpropyl)-5-(4-fluoro-2-methoxyphenyl)pyrazin-2-amine
192		6-ethyl-N-(1-ethylpropyl)-5-(4-fluoro-2-methoxyphenyl)-3-methoxypyrazin-2-amine
193		3-ethyl-N-(1-ethylpropyl)-5-(4-fluoro-2-methoxyphenyl)-6-methoxypyrazin-2-amine
194		3-ethyl-N-(1-ethylpropyl)-6-methoxy-5-[4-(trifluoromethyl)phenyl]pyrazin-2-amine







195		5-(2-chloro-4-methylphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
196		2,5-diethyl-3-(1-ethylbutoxy)-6-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazine
197		2,5-diethyl-3-(1-ethylpropoxy)-6-[2-methoxy-4-(trifluoromethyl)phenyl]pyrazine
198		5-(2-chloro-4-methylphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
199		5-(2-chloro-4-methylphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
200		5-(4-chloro-2-methoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine

201		5-(4-chloro-2-methoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
202		5-(4-chloro-2-methoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
203		5-(2-chloro-4-isopropoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
204		O-(3-chloro-4-{3,6-diethyl-5-[(1-ethylpropyl)amino]pyrazin-2-yl}phenyl) S-propyl (dithiocarbonate)
205		5-(2,6-dimethylphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
206		5-(2,6-dimethoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







207		5-(4-chloro-2,6-dimethoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
208		5-(4-chloro-2,6-dimethoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
209		5-(3,5-dichlorothiophen-2-yl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
210		5-(3,5-dichlorothiophen-2-yl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
211		2-(2,4-dimethoxyphenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
212		2-(cyclopentyloxy)-5-(2,4-dimethoxyphenyl)-3,6-diethylpyrazine

213		2,5-diethyl-3-(1-ethylbutoxy)-6-[2-methoxy-4-(trifluoromethyl)phenyl]pyrazine
214		2,5-diethyl-3-(1-ethylbutoxy)-6-[4-methoxy-2-(trifluoromethyl)phenyl]pyrazine
215		5-(4-tert-butyl-2,6-dimethylphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
216		5-(2,6-dichloro-4-methoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
217		5-[2-chloro-4-(difluoromethoxy)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine
218		5-(2,6-dimethoxy-4-methylphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







219		2-(2-chloro-4-methoxyphenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
220		5-[4-(1,3-dioxolan-2-yl)-2,6-dimethoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
221		N-(1-ethylpropyl)-5-(4-fluoro-2-methoxyphenyl)-3-methoxy-6-methylpyrazin-2-amine
222		5-(2-chloro-4-fluorophenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
223		5-[2,4-bis(trifluoromethyl)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
224		5-(4-chloro-2-methylphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







225		5-(4-chloro-2-methoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
226		5-(2,4-diethoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
227		5-(4-ethoxy-2-methoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
228		N-(1-ethylpropyl)-3-methoxy-5-(2-methoxy-4-methylphenyl)-6-methylpyrazin-2-amine
229		3-ethyl-N-(1-ethylpropyl)-6-methoxy-5-(2-methoxy-6-methylphenyl)pyrazin-2-amine
230		3-ethyl-N-(1-ethylpropyl)-6-methoxy-5-[4-methoxy-2-(trifluoromethyl)phenyl]pyrazin-2-amine







231		6-ethyl-N-(1-ethylpropyl)-3-methoxy-5-(2-methoxy-6-methylphenyl)pyrazin-2-amine
232		N-(1-ethylpropyl)-3-methoxy-5-(2-methoxy-6-methylphenyl)pyrazin-2-amine
233		5-[4-(difluoromethyl)-2,6-dimethoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
234		1-(4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-3,5-dimethoxyphenyl)ethanol
235		1-(4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-3,5-dimethoxyphenyl)ethanone
236		5-(2-chloro-4-methoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







237		2-[2,4-bis(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
238		2-(2-chloro-4-methoxyphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
239		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[2-methoxy-4-(trifluoromethyl)phenyl]pyrazine
240		5-(2-chloro-4-ethoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
241		5-(2-chloro-4-isopropoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
242		5-(2-chloro-4-isopropoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine







243		2-(4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-3,5-dimethoxyphenyl)propan-2-ol
244		2,5-diethyl-3-(1-ethylbutoxy)-6-(2-methoxy-4,6-dimethylphenyl)pyrazine
245		2,5-diethyl-3-(1-ethylbutoxy)-6-[4-methoxy-2-(trifluoromethoxy)phenyl]pyrazine
246		5-(4-chloro-2-ethoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
247		5-(4-chloro-2-ethoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
248		3-ethyl-6-methoxy-N-(4-methoxybenzyl)-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine







249		5-(2,4-dimethoxyphenyl)-3-ethyl-6-methoxy-N-(4-methoxybenzyl)pyrazin-2-amine
250		2-(2,4-dichloro-6-methylphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
251		2-(4-chloro-2,6-dimethoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
252		3-ethyl-5-(2-ethyl-4-methoxyphenyl)-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
253		5-(2-ethyl-4-methoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
254		6-ethyl-5-(2-ethyl-4-methoxyphenyl)-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine

255		5-[4-(benzyloxy)-2-isopropoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
256		5-(4-chloro-2-ethoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
257		N-(1-ethylpropyl)-3-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
258		3-chloro-4-{6-ethyl-5-[(1-ethylpropyl)amino]-3-methoxypyrazin-2-yl}phenol
259		3-chloro-4-{3-ethyl-5-[(1-ethylpropyl)amino]-6-methoxypyrazin-2-yl}phenol
260		5-(2-chloro-4-isopropoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine

261		5-(2-chloro-4-methoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
262		5-[2-chloro-4-(difluoromethoxy)phenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
263		5-[2-chloro-4-(difluoromethoxy)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
264		5-[2-chloro-4-(1-ethylpropoxy)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
265		5-[2-chloro-4-(2-fluoroethoxy)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
266		5-[2-chloro-4-(difluoromethoxy)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







267		5-[2-chloro-4-(1-ethylpropoxy)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
268		3-(benzyloxy)-4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}phenol
269		5-[4-(but-3-enyloxy)-2-chlorophenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
270		5-(2,4-dimethoxyphenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
271		5-(2,4-dichlorophenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
272		N-(1-ethylpropyl)-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]-3-methylpyrazin-2-amine







273		5-[2-chloro-4-(trifluoromethyl)phenyl]-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
274		5-[2'-chloro-4',5-bis(trifluoromethyl)-1,1'-biphenyl-2-yl]-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
275		N-(1-ethylpropyl)-6-methoxy-5-[2-methoxy-4-(trifluoromethyl)phenyl]-3-methylpyrazin-2-amine
276		5-(2-chloro-4-methylphenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
277		N-(1-ethylpropyl)-6-methoxy-3-methyl-5-(4-methylphenyl)pyrazin-2-amine
278		N-(1-ethylpropyl)-5-(4-fluoro-2-methoxyphenyl)-6-methoxy-3-methylpyrazin-2-amine

279		N-(1-ethylpropyl)-6-methoxy-3-methyl-5-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
280		5-(2-ethyl-4-methoxyphenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
281		N-(1-ethylpropyl)-6-methoxy-3-methyl-5-[4-methyl-2-(trifluoromethyl)phenyl]pyrazin-2-amine
282		5-(2,4-dimethylphenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
283		5-(4-chloro-2,6-dimethoxyphenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
284		5-(4-chloro-2,6-dimethoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine







285		2,5-diethyl-3-(2-ethyl-4-methoxyphenyl)-6-(1-isopropyl-2-methylpropoxy)pyrazine
286		2-(4-chloro-2,6-dimethoxyphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
287		4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-3-isopropoxyphenol
288		5-(2,4-diisopropoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
289		N-(1-ethylpropyl)-5-(2-isopropoxy-4-propoxyphenyl)-3-methoxy-6-methylpyrazin-2-amine
290		5-(4-ethoxy-2-isopropoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







291		(3-chloro-4-((1-ethylpropyl)amino)-6-methoxy-3-methylpyrazin-2-yl)phenoxy)acetonitrile
292		5-(2-chloro-4-fluorophenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine
293		5-[2-chloro-4-(1-ethylpropoxy)phenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
294		5-[4-(cyclopentyloxy)-2-isopropoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
295		5-[4-(allyloxy)-2-isopropoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
296		2-[2-chloro-4-(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-ethylbutoxy)pyrazine







297		2-[2-chloro-4-(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
298		2-(4-chloro-2-methylphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
299		N-(1-ethylpropyl)-5-[4-(1-fluoroethyl)-2,6-dimethoxyphenyl]-3-methoxy-6-methylpyrazin-2-amine
300		2-(2-chloro-4-fluorophenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
301		5-(2-chloro-4-[(Z)-1-fluoro-2-iodoethenyl]oxy)phenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
302		5-[2-chloro-4-(2-methoxyethoxy)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







303		5-(4-((tert-butyl(dimethyl)silyloxy)-2-methoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
304		4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-3-methoxyphenol
305		5-[4-(difluoromethoxy)-2-methoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine
306		2-(2-chloro-4-fluorophenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
307		2-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-5-methoxybenzaldehyde
308		5-{2-[(dimethylamino)methyl]-4-methoxyphenyl}-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







309		N-(1-ethylpropyl)-3-methoxy-5-[4-methoxy-2-(morpholin-4-ylmethyl)phenyl]-6-methylpyrazin-2-amine
310		N-(1-ethylpropyl)-3-methoxy-5-[4-methoxy-2-(piperidin-1-ylmethyl)phenyl]-6-methylpyrazin-2-amine
311		(2-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-5-methoxyphenyl)methanol
312		1-(2-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-5-methoxyphenyl)ethanol
313		1-(2-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-5-methoxyphenyl)ethanone
314		5-[4-(difluoromethoxy)-2-methoxyphenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine







315		3-chloro-4-[3,6-diethyl-5-(1-ethylpropoxy)pyrazin-2-yl]phenol
316		2-(2-chloro-4-ethoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
317		2-(2-chloro-4-propoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
318		2-(2-chloro-4-isopropoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
319		2-(2-chloro-4-isobutoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
320		2-[2-chloro-4-(cyclopropylmethoxy)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine







321		2-(4-butoxy-2-chlorophenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
322		{3-chloro-4-[3,6-diethyl-5-(1-ethylpropoxy)pyrazin-2-yl]phenoxy} acetonitrile
323		2-[2-chloro-4-(2-fluoroethoxy)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
324		2-[2-chloro-4-(2-methoxyethoxy)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
325		4-{6-ethyl-5-[(1-ethylpropyl)amino]-3-methoxypyrazin-2-yl}-3-methoxyphenol
326		5-(4-ethoxy-2-methoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine







327		N-(1-ethylpropyl)-5-(4-isopropoxy-2-methoxyphenyl)-3-methoxy-6-methylpyrazin-2-amine
328		3-ethyl-N-(1-ethylpropyl)-5-(4-isopropoxy-2-methoxyphenyl)-6-methoxypyrazin-2-amine
329		2-(4-{{tert-butyl(dimethyl)silyl}oxy}-2-isopropoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
330		2-[2-chloro-4-(difluoromethoxy)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
331		2-[4-(difluoromethoxy)-2-methoxyphenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
332		4-[3,6-diethyl-5-(1-ethylpropoxy)pyrazin-2-yl]-3-methoxyphenol







333		2-(4-ethoxy-2-methoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
334		2,5-diethyl-3-(1-ethylpropoxy)-6-(4-isopropoxy-2-methoxyphenyl)pyrazine
335		2,5-diethyl-3-(1-ethylbutoxy)-6-(4-fluoro-2-methylphenyl)pyrazine
336		2,5-diethyl-3-(4-fluoro-2-methylphenyl)-6-(1-isopropyl-2-methylpropoxy)pyrazine
337		2-(2,6-dimethoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
338		2-(2,6-dimethoxyphenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine







339		2-(2,6-dimethoxyphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
340		4-[3,6-diethyl-5-(1-ethylpropoxy)pyrazin-2-yl]-3-isopropoxyphenol
341		4-{3-ethyl-5-[(1-ethylpropyl)amino]-6-methoxypyrazin-2-yl}-3-methoxyphenol
342		5-[4-(difluoromethoxy)-2-methoxyphenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
343		5-(4-ethoxy-2-methoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine
344		6-ethyl-N-(1-ethylpropyl)-5-(4-isopropoxy-2-methoxyphenyl)-3-methoxypyrazin-2-amine







345		2-(2,4-diisopropoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
346		N-cyclopentyl-5-(2,4-dichlorophenyl)-3,6-diethylpyrazin-2-amine
347		2,5-diethyl-3-(2-ethyl-4-methoxyphenyl)-6-(1-ethylpropoxy)pyrazine
348		4-[3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazin-2-yl]-3-ethylphenol
349		2-(4-ethoxy-2-ethylphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
350		2,5-diethyl-3-(2-ethyl-4-isopropoxyphenyl)-6-(1-isopropyl-2-methylpropoxy)pyrazine

351		2-[4-(cyclopentyloxy)-2-ethylphenyl]-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
352		2-(4-ethoxy-2-isopropoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
353		2-(2,4-dimethoxyphenyl)-3,6-diethyl-5-(1-propylbutoxy)pyrazine
354		2-(2-chloro-4-methoxyphenyl)-3,6-diethyl-5-(1-propylbutoxy)pyrazine
355		3-ethyl-N-(1-ethylpropyl)-5-[4-fluoro-2-(trifluoromethyl)phenyl]-6-methoxypyrazin-2-amine
356		3-chloro-4-{3,6-diethyl-5-[(1-ethylpropyl)amino]pyrazin-2-yl}phenol







357		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[2-methoxy-4,6-bis(trifluoromethyl)phenyl]pyrazine
358		2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-mesitylpyrazine
359		2-(2,4-dimethylphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine
360		2,5-diethyl-3-(1-ethylpropoxy)-6-mesitylpyrazine
361		2,5-diethyl-3-(1-ethylpropoxy)-6-[4-methoxy-2-(trifluoromethyl)phenyl]pyrazine
362		2,5-diethyl-3-(1-ethylpropoxy)-6-(4-methoxy-2-methylphenyl)pyrazine





363		5-[2,4-bis(trifluoromethyl)phenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
364		5-(4-chloro-2-methylphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine
365		2,5-diethyl-3-[2-methoxy-4-(trifluoromethoxy)phenyl]-6-(1-propylbutoxy)pyrazine
366		2,5-diethyl-3-[4-methoxy-2-(trifluoromethoxy)phenyl]-6-(1-propylbutoxy)pyrazine
367		2,5-diethyl-3-(4-fluoro-2-methylphenyl)-6-(1-propylbutoxy)pyrazine
368		2,5-diethyl-3-[2-methoxy-4-(trifluoromethyl)phenyl]-6-(1-propylbutoxy)pyrazine

369		6-chloro-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
370		2-(2,4-dimethoxyphenyl)-3,6-diethyl-5-(2-methylbutyl)pyrazine
371		2-(2-chloro-4-ethoxyphenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
372		2-(2-chloro-4-isopropoxyphenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
373		2-(4-butoxy-2-chlorophenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
374		2-[2-chloro-4-(cyclopentyloxy)phenyl]-3,6-diethyl-5-(1-ethylbutoxy)pyrazine

375		2-[2-chloro-4-(difluoromethoxy)phenyl]-3,6-diethyl-5-(1-ethylbutoxy)pyrazine
376		N-(1-ethylpropyl)-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
377		3-bromo-N-(1-ethylpropyl)-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
378		3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine
379		2-ethyl-3-(1-ethylpropoxy)-5-methoxy-6-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazine
380		2-(4-[[tert-butyl(dimethyl)silyl]oxy]-2-ethoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine

381		4-[3,6-diethyl-5-(1-ethylpropoxy)pyrazin-2-yl]-3-ethoxyphenol
382		2-(2,4-diethoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
383		2-(2-ethoxy-4-isopropoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
384		6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-ylformamide
385		3-ethyl-N-(1-ethylpropyl)-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine 4-oxide
386		6-chloro-3-ethyl-N-(1-ethylpropyl)-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine

387		2,5-diethyl-3-(2-ethylbutyl)-6-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazine
388		2-[2-chloro-4-(trifluoromethyl)phenyl]-3,6-diethyl-5-(2-ethylbutyl)pyrazine
389		2-(2-ethoxy-4-propoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine
390		2,5-diethyl-3-(1-ethylpropoxy)-6-(2-isopropoxy-4-propoxyphenyl)pyrazine
391		N-[6-chloro-5-(4-chloro-2,6-dimethoxyphenyl)pyrazin-2-yl]acetamide
392		2-(sec-butylthio)-3-ethyl-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazine


393		2-(sec-butylthio)-5-(3,6-dichlorophenyl)-3,6-diethylpyrazine
394		2-(sec-butylthio)-5-(2,4-dimethoxyphenyl)-3,6-diethylpyrazine
395		3-ethyl-N-(1-ethylpropyl)-5-[2-methoxy-4-(trifluoromethoxy)phenyl]-6-(methylthio)pyrazin-2-amine
396		5-(4-chloro-2,6-dimethoxyphenyl)-6-methoxypyrazin-2-amine

The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the present invention and that

5 modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

WHAT IS CLAIMED IS:

1. A compound of the formula:



5

or a pharmaceutically acceptable salt thereof, wherein:

Ar is substituted phenyl, optionally substituted naphthyl, or an optionally substituted heterocyclic group having from 1 to 3 rings, and 3 to 8 ring members in each ring and 1 to about 3 hetero atoms;

10 R₁ and R₃ are each independently hydrogen, halogen, cyano, nitro, amino, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted mono or dialkylamino, optionally substituted alkoxy, optionally substituted alkylthio, optionally substituted alkylsulfinyl, or optionally substituted alkylsulfonyl; and


15 R₂ is halogen, cyano, nitro, amino, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkylamino, optionally substituted alkoxy, optionally substituted alkylthio, optionally substituted alkylsulfinyl, or optionally substituted alkylsulfonyl;

20 with the proviso that if Ar is phenyl substituted with halogen, naphthyl, or naphthyl substituted with halogen, then the compounds where R₃ is hydrogen or amino are excluded.

2. A compound according to Claim 1, wherein:

25 Ar is substituted phenyl, optionally substituted naphthyl, or an optionally substituted heterocyclic group having at least one nitrogen ring atom or at least one sulfur ring atom.

3. A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

R₁ is selected from H, C₁₋₄ alkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, halogen, CN, C₁₋₄ haloalkyl, trifluoromethyl, trifluoromethoxy, -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), -O(C₁₋₄ alkyl), and S(O)_n(C₁₋₄ alkyl);

R₂ is selected from the group consisting of -XR_A and Y, wherein -X, R_A, and Y are defined below; and

R₃ is selected from the group consisting of hydrogen, halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), and -S(O)_n(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, trifluoromethoxy, -XR_A and Y;

R₄ is absent or an oxygen atom;

Ar is phenyl, mono-, di-, or tri-substituted with R_C, or

Ar is selected from the group consisting of:

naphthyl, pyridyl, pyridonyl, pyrimidinyl, and thiophenyl, each of which is unsubstituted or mono-, di-, or tri-substituted with R_C;

with the proviso that if Ar is phenyl substituted with halogen, naphthyl, or naphthyl substituted with halogen, then the compounds where R₃ is hydrogen are excluded;

R_A and R_B, which may be the same or different, are independently selected at each occurrence from the group consisting of: hydrogen and straight, branched, or cyclic alkyl groups consisting of 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) selected from oxo, hydroxy, halogen, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), -NHC(O)(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)C(=O)(C₁₋₄ alkyl), -NHS(O)_n(C₁₋₄ alkyl), -S(O)_n(C₁₋₄ alkyl), -S(O)_nNH(C₁₋₄ alkyl), -S(O)_nN(C₁₋₄ alkyl)(C₁₋₄ alkyl), and Z;

R_C is independently selected at each occurrence from the group consisting of halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆

alkyl optionally substituted with 0-2 R_D , C_{2-6} alkenyl substituted with 0-2 R_D , C_{1-4} alkynyl substituted with 0-2 R_D , C_{3-7} cycloalkyl substituted with 0-2 R_D , (C_{3-7} cycloalkyl) C_{1-4} alkyl substituted with 0-2 R_D , $-O(C_{1-4}$ alkyl) substituted with 0-2 R_D , $-NH(C_{1-4}$ alkyl) substituted with 0-2 R_D , $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl) each
 5 independently substituted with 0-2 R_D , $-XR_A$, and Y;

R_D is independently selected at each occurrence the group consisting of halogen, hydroxy, cyano, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), morpholino, pyrrolidino, piperidino, thiomorpholino, piperazino, 4-hydroxypiperidino, $-S(O)_n(C_{1-4}$ alkyl), trifluoromethyl, trifluoromethoxy, $CO(C_{1-4}$ alkyl), $CONH(C_{1-4}$ alkyl), $CON(C_{1-4}$ alkyl)(C_{1-4} alkyl), $-XR_A$ and Y;
 10

X is independently selected at each occurrence from the group consisting of $-CH_2-$, $-CHR_B-$, $-O-$, $-C(=O)-$, $-C(=O)O-$, $-S(O)_n-$, $-NH-$, $-NR_B-$, $-C(=O)NH-$, $-C(=O)NR_B-$, $-S(O)_nNH-$, $-S(O)_nNR_B-$, $-OC(=S)S-$, $-NHC(=O)-$, $-NR_BC(=O)-$, $-NHS(O)_n-$, $-OSiH_n(C_{1-4}alkyl)_{2-n}-$, and $-NR_BS(O)_n-$; and

15 Y and Z are independently selected at each occurrence from the group consisting of: 3- to 7-membered carbocyclic and heterocyclic groups, which are saturated, unsaturated, or aromatic, which may be substituted with one or more substituents selected from halogen, haloalkyl, oxo, hydroxy, amino, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), and $-S(O)_n(C_{1-4}$ alkyl), and
 20 said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) selected from N, O, and S, with the point of attachment being either carbon or nitrogen; and

n is independently selected at each occurrence from 0, 1, and 2.

25 4. A compound of according to Claim 1 wherein Ar is substituted phenyl.

5. A compound according to Claim 3, R_4 is absent and Ar is phenyl, mono-, di-, or tri-substituted with R_C .


6. A compound according to Claim 3 wherein R_4 is absent and Ar is phenyl, mono-, di-, or tri-substituted with R_C , and R_1 and R_3 are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy and methoxy.

7. A compound according to Claim 3 wherein R_4 is absent and Ar is phenyl, mono-, di-, or tri-substituted with R_C ; and R_A and R_B , which may be the same or different, are independently straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds.

8. A compound according to Claim 3 wherein R_4 is absent and Ar is phenyl mono-, di-, or tri-substituted with R_C ; R_A and R_B , which may be the same or different, are independently selected at each occurrence from the group consisting of: straight, branched, and cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds; and

R_1 and R_3 are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy, and methoxy.

9. A compound of the Formula:



Formula A

wherein

R_X and R_Y are the same or different and are independently selected straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) independently selected from hydroxy, halogen, $-O(C_{1-4} \text{ alkyl})$, $-NH(C_{1-4} \text{ alkyl})$, $-N(C_{1-4} \text{ alkyl})(C_{1-4} \text{ alkyl})$, and optionally substituted phenyl.

10. A compound according to Claim 9, wherein Ar is phenyl, mono-, di-, or tri-substituted with R_C ; and R_1 and R_3 are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy and methoxy.

5

11. A compound according to Claim 9, wherein:

R_1 and R_3 are independently chosen from halogen, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

10 Ar is phenyl, which is mono-, di-, or trisubstituted with one or more substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy(C_{1-4} alkoxy), mono- or di(C_{1-4})amino(C_{1-4} alkoxy), and mono- or di(C_{1-4} alkyl)amino.

15

12. A compound according to Claim 9, wherein:

R_X is hydrogen;

R_Y is chosen from the group consisting of:

straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may
20 contain one or more double or triple bonds;

R_1 and R_3 are independently chosen from halogen, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

25 Ar is phenyl, which is mono-, di-, or trisubstituted with substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy(C_{1-4} alkoxy), mono- or di(C_{1-4})amino(C_{1-4} alkoxy), and mono- or di(C_{1-4} alkyl)amino.

30

13. A compound according to Claim 9, wherein:


R_x is hydrogen;

R_y is chosen from the group consisting of:

straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds;

- 5 R_1 and R_3 are independently chosen from halogen, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

Ar is a phenyl group of the formula:



- 10 wherein L indicates a bond to the pyrazine ring in Formula A;
and the phenyl group is substituted at one, two or three of positions 2, 4, and 6 with substituent(s) independently selected from:
halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino,
and C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy(C_{1-4} alkoxy), mono- or di(C_{1-4})amino(C_{1-4} alkoxy), and mono- or di(C_{1-4} alkyl)amino.
- 15

14. A compound according to Claim 9, wherein:


R_x is hydrogen;

R_y is chosen from the group consisting of:

- 20 straight, branched, or cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds;

R_1 and R_3 are independently chosen from halogen, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

- 25 Ar is a phenyl group of the formula:




wherein L indicates a bond to the pyrazine ring in Formula A;

and the phenyl group is substituted at positions 2 and 4 with substituents independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.


15. A compound according to Claim 9 of the formula:



wherein:

R₁ and R₃ are independently chosen at each occurrence from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

Ar is a phenyl group of the formula:




wherein L indicates a bond to the pyrazine ring in Formula A;

and the phenyl group is substituted at one, two or three of positions 2, 4, and 6 with substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.

16. A compound according to Claim 9 of the formula:



wherein

R is independently selected at each occurrence from the group consisting of: hydrogen, halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, C₁₋₆alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄alkyl)amino; and


R₁ and R₃ are independently chosen from halogen, C₁₋₄alkyl, -O(C₁₋₄alkyl), -NH(C₁₋₄alkyl), -N(C₁₋₄alkyl)(C₁₋₄alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy.

17. A compound according to Claim 9 wherein Ar is phenyl, which is mono-, di-, or tri-substituted with R_C;

R_X and R_Y, which may be the same or different, are independently selected at each occurrence from the group consisting of: straight, branched, and cyclic alkyl groups having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds; and


R₁ and R₃ are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy, and methoxy.

18. A compound according to Claim 3 of the formula:




wherein R_X and R_Y are independently hydrogen or C₁₋₈alkyl; or

NR_xR_y is:



5 wherein z is 0 or 1.

19. A compound of the Formula:



10 Formula B

wherein:

R_x is chosen from the group consisting of:

straight, branched, or cyclic alkyl groups, including (cycloalkyl)alkyl groups, having from 1 to 8 carbon atoms, which may contain one or more double or triple bonds, each of which may be further substituted with one or more substituent(s) independently selected from

- 15
- (a) hydroxy, halogen, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), and
 - (b) 3- to 7-membered carbocyclic and heterocyclic groups, which are
- 20
- saturated, unsaturated, or aromatic, which may be substituted with one or more substituents selected from halogen, haloalkyl, oxo, hydroxy, amino, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), and -N(C₁₋₄ alkyl)(C₁₋₄ alkyl) and wherein said 3- to 7-membered heterocyclic groups contain one or more heteroatom(s) selected from N, O, and S, with the point of attachment
- 25
- being either carbon or nitrogen.

20. A compound according to Claim 19, wherein:

R_x is selected from straight, branched, or cyclic alkyl groups containing of 1 to 8 carbon atoms, which may contain one or more double or triple bonds;

5 and;


R_1 and R_3 are independently chosen from halogen, C_{1-4} alkyl, $-O(C_{1-4}$ alkyl), $-NH(C_{1-4}$ alkyl), $-N(C_{1-4}$ alkyl)(C_{1-4} alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and

10 Ar is phenyl, which is mono-, di-, or trisubstituted with one or more substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy(C_{1-4} alkoxy), mono- or di(C_{1-4})amino(C_{1-4} alkoxy), and mono- or di(C_{1-4} alkyl)amino.

15 21. A compound according to Claim 20, wherein:

Ar is a phenyl group of the formula:




wherein L indicates a bond to the pyrazine ring in Formula B;

20 and the phenyl group is substituted at one, two, or three of positions 2, 4, and 6 with substituent(s) independently selected from:

halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, and C_{1-6} alkyl, C_{1-4} alkoxy, C_{1-4} alkoxy(C_{1-4} alkoxy), mono- or di(C_{1-4})amino(C_{1-4} alkoxy), and mono- or di(C_{1-4} alkyl)amino.

25 22. A compound according to Claim 20, wherein:


Ar is a phenyl group of the formula:



wherein L indicates a bond to the pyrazine ring in Formula B;

and the phenyl group is substituted at positions 2 and 4 with substituents independently selected from halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.


23. A compound according to Claim 19, of the formula:



wherein:

- 10 R₁ and R₃ are independently chosen from halogen, C₁₋₄ alkyl, -O(C₁₋₄ alkyl), -NH(C₁₋₄ alkyl), -N(C₁₋₄ alkyl)(C₁₋₄ alkyl), haloalkyl, trifluoromethyl, and trifluoromethoxy; and


Ar is a phenyl group of the formula:



- 15 wherein L indicates a bond to the pyrazine ring in Formula B;
and the phenyl group is substituted at one, two, or three of positions 2, 4, and 6 with substituent(s) independently selected from: halogen, cyano, haloalkyl, trifluoromethyl, trifluoromethoxy, hydroxy, amino, C₁₋₆ alkyl, C₁₋₄alkoxy, C₁₋₄alkoxy(C₁₋₄alkoxy), mono- or di(C₁₋₄)amino(C₁₋₄alkoxy), and mono- or di(C₁₋₄ alkyl)amino.
- 20

24. A compound according to Claim 19 wherein R₁ and R₃ are independently selected from the group consisting of hydrogen, halogen, methyl, ethyl, ethoxy and methoxy.

25. A compound according to Claim 3 of the formula:



5 wherein A is NR_A or O.

26. A compound according to Claim 1 wherein, in a standard in vitro CRF receptor binding assay the compound exhibits an IC₅₀ value less than or equal to 1 micromolar.

10

27. A method for modulating CRF receptor function, comprising administering to a patient in need of such modulation an effective amount of a compound of according to Claim 1.

15

28. A method for treating a CNS disorder or disease, comprising administering to a patient in need of such treatment an effective amount of a compound according to Claim 1.

20

29. The method of Claim 28 wherein the CNS disorder or disease is an anxiety disorder, stress-related disorder, or eating disorder.

30. A compound according to Claim 1, wherein in a standard in vitro sodium channel functional assay the compound does not show any statistically significant activity at the $p < 0.05$ level of significance.

25

31. A method of Claim 28 wherein the CNS disease or disorder is depression or a bipolar disorder.

32. A method of Claim 28 wherein the CNS disease or disorder is anorexia nervosa, bulimia nervosa, or obesity.

5 33. A method according to Claim 28 wherein in a standard in vitro CRF receptor binding assay the compound exhibits an IC_{50} value less than or equal to 1 micromolar.

34. A method according to Claim 28 wherein in a standard in vitro CRF
10 receptor binding assay the compound exhibits an IC_{50} value less than or equal to 100 nanomolar.

35. A method according to Claim 28 wherein in a standard in vitro CRF
15 receptor binding assay the compound exhibits an IC_{50} value less than or equal to 10 nanomolar.

36. A method according to Claim 28 wherein the patient is a human.

37. A method for localizing CRF receptors is tissue section samples
20 comprising:

contacting with a sample of tissue a detectably-labelled compound of Claim 1
under conditions that permit binding of the compound to CRF receptors within the
sample of tissue;

washing the tissue sample to remove unbound compound; and
25 detecting remaining bound compound.

38. A method of inhibiting the binding of CRF to a CRF1 Receptor which
comprises:

contacting a solution comprising CRF and a compound of Claim 3 with a cell the receptor, wherein the compound is present in the solution at a concentration sufficient to inhibit CRF binding to IMR32 cells in vitro.

5 39. A method for altering the signal-transducing activity of a CRF1 receptor, the method comprising contacting cells expressing the receptor with a compound according to Claim 3.

10 40. A method of modulating G-protein coupled receptor function, the method comprising exposing cells expressing a G-protein coupled receptor to an effective amount of a compound according to Claim 3.

15 41. A method according to Claim 40, wherein the G-coupled protein receptor is a CRF1 receptor.

 42. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of a compound according to Claim 1.

20 43. A packaged pharmaceutical composition comprising a pharmaceutical composition of claim 42 in a container and instructions for using the composition to treat a patient suffering from an anxiety disorder, a stress-related disorder, or an eating disorder.

25 44. A compound according to Claim 3, named 5-(2,4-dimethoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

 45. A compound according to Claim 3, named 5-[4-bromo-2-(trifluoromethoxy)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine.

30 46. A compound according to Claim 3, named 3-ethyl-N-(1-ethylpropyl)-6-

methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl] pyrazin-2-amine.

47. A compound according to Claim 3, named 5-(4-chloro-2-methylphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

5

48. A compound according to Claim 3, named 5-[2-chloro-4-(trifluoromethyl)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

49. A compound according to Claim 3, named 2-[2,4-bis(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine.

10

50. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(2-naphthyl)pyrazin-2-amine.

51. A compound according to Claim 3, named 5-[4-chloro-2-(trifluoromethyl)phenyl]-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine.

15

52. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-[2-methoxy-6-(trifluoromethoxy)phenyl]pyrazin-2-amine.

20

53. A compound according to Claim 3, named 5-(2-chloro-4-methoxyphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

54. A compound according to Claim 3, named 5-(2,6-dichloro-4-methoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine.

25

55. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(3-methyl-1,1'-biphenyl-4-yl)pyrazin-2-amine.

56. A compound according to Claim 3, named 5-[2,4-

30

bis(trifluoromethyl)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

57. A compound according to Claim 3, named 2-(2,4-dichlorophenyl)-3,6-diethyl-5-(1-methylbutoxy)pyrazine.

5

58. A compound according to Claim 3, named 2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[4-methoxy-2-(trifluoromethoxy)phenyl]pyrazine.

59. A compound according to Claim 3, named 2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazine.

10

60. A compound according to Claim 3, named 5-[4-chloro-2-(trifluoromethyl)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

61. A compound according to Claim 3, named N-(1-ethylpropyl)-3-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]-6-methylpyrazin-2-amine.

15

62. A compound according to Claim 3, named 5-[4-chloro-2-(trifluoromethyl)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

20

63. A compound according to Claim 3, named 3-ethyl-N-(1-ethylpropyl)-6-methoxy-5-[2-methyl-4-(trifluoromethoxy)phenyl]pyrazin-2-amine.

64. A compound according to Claim 3, named 5-(2,4-dimethylphenyl)-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

25

65. A compound according to Claim 3, named 5-(2-chloro-4-ethoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

66. A compound according to Claim 3, named O-(3-chloro-4-{5-[(1-

30

ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}phenyl) S-propyl (dithiocarbonate).

67. A compound according to Claim 3, named 2,5-diethyl-3-(1-ethylpropoxy)-6-[2-methoxy-4,6-bis(trifluoromethyl)phenyl]pyrazine.

5

68. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(4-fluoro-2-methoxyphenyl)pyrazin-2-amine.

69. A compound according to Claim 3, named 5-(2-chloro-4-methylphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine.

10

70. A compound according to Claim 3, named 5-(4-chloro-2-methoxyphenyl)-3,6-diethyl-N-(1-ethylpropyl)pyrazin-2-amine.

71. A compound according to Claim 3, named O-(3-chloro-4-{3,6-diethyl-5-[(1-ethylpropyl)amino]pyrazin-2-yl}phenyl) S-propyl (dithiocarbonate).

15

72. A compound according to Claim 3, named 5-(4-chloro-2,6-dimethoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

20

73. A compound according to Claim 3, named 2-(2,4-dimethoxyphenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine.

74. A compound according to Claim 3, named 5-[4-(1,3-dioxolan-2-yl)-2,6-dimethoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

25

75. A compound according to Claim 3, named 5-(4-ethoxy-2-methoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine.

76. A compound according to Claim 3, named 6-ethyl-N-(1-ethylpropyl)-3-

30

methoxy-5-(2-methoxy-6-methylphenyl)pyrazin-2-amine.

77. A compound according to Claim 3, named 2-(2-chloro-4-methoxyphenyl)-3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazine.

5

78. A compound according to Claim 3, named 2-(4-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-3,5-dimethoxyphenyl)propan-2-ol.

79. A compound according to Claim 3, named 5-(4-chloro-2-ethoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

10

80. A compound according to Claim 3, named 3-ethyl-5-(2-ethyl-4-methoxyphenyl)-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine.

81. A compound according to Claim 3, named 5-(4-chloro-2-ethoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine.

15

82. A compound according to Claim 3, named 5-(2-chloro-4-isopropoxyphenyl)-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine.

20

83. A compound according to Claim 3, named 5-[2-chloro-4-(1-ethylpropoxy)phenyl]-6-ethyl-N-(1-ethylpropyl)-3-methoxypyrazin-2-amine.

84. A compound according to Claim 3, named 5-[2-chloro-4-(1-ethylpropoxy)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

25

85. A compound according to Claim 3, named 5-(2,4-dichlorophenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine.

86. A compound according to Claim 3, named 5-[2'-chloro-4',5-

30

bis(trifluoromethyl)-1,1'-biphenyl-2-yl]-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine.

87. A compound according to Claim 3, named N-(1-ethylpropyl)-5-(4-fluoro-2-methoxyphenyl)-6-methoxy-3-methylpyrazin-2-amine.

88. A compound according to Claim 3, named 5-(2,4-dimethylphenyl)-N-(1-ethylpropyl)-6-methoxy-3-methylpyrazin-2-amine.

89. A compound according to Claim 3, named 2,5-diethyl-3-(2-ethyl-4-methoxyphenyl)-6-(1-isopropyl-2-methylpropoxy)pyrazine.

90. A compound according to Claim 3, named 5-(4-ethoxy-2-isopropoxyphenyl)-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

91. A compound according to Claim 3, named 5-[2-chloro-4-(1-ethylpropoxy)phenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine.

92. A compound according to Claim 3, named 2-[2-chloro-4-(trifluoromethyl)phenyl]-3,6-diethyl-5-(1-ethylbutoxy)pyrazine.

93. A compound according to Claim 3, named N-(1-ethylpropyl)-5-[4-(1-fluoroethyl)-2,6-dimethoxyphenyl]-3-methoxy-6-methylpyrazin-2-amine.

94. A compound according to Claim 3, named 5-[2-chloro-4-(2-methoxyethoxy)phenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

95. A compound according to Claim 3, named 5-[4-(difluoromethoxy)-2-methoxyphenyl]-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

96. A compound according to Claim 3, named 5-{2-[(dimethylamino)methyl]-4-methoxyphenyl}-N-(1-ethylpropyl)-3-methoxy-6-methylpyrazin-2-amine.

97. A compound according to Claim 3, named (2-{5-[(1-ethylpropyl)amino]-6-methoxy-3-methylpyrazin-2-yl}-5-methoxyphenyl)methanol.

98. A compound according to Claim 3, named 5-[4-(difluoromethoxy)-2-methoxyphenyl]-3-ethyl-N-(1-ethylpropyl)-6-methoxypyrazin-2-amine.

99. A compound according to Claim 3, named 2-(2-chloro-4-propoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine.

100. A compound according to Claim 3, named 2-[2-chloro-4-(cyclopropylmethoxy)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine.

101. A compound according to Claim 3, named 2-[2-chloro-4-(2-fluoroethoxy)phenyl]-3,6-diethyl-5-(1-ethylpropoxy)pyrazine.

102. A compound according to Claim 3, named N-(1-ethylpropyl)-5-(4-isopropoxy-2-methoxyphenyl)-3-methoxy-6-methylpyrazin-2-amine.

103. A compound according to Claim 3, named 2-(4-[[tert-butyl(dimethyl)silyl]oxy]-2-isopropoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine.

104. A compound according to Claim 3, named 2-(4-ethoxy-2-methoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine.

105. A compound according to Claim 3, named 2-(2,6-dimethoxyphenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine.

106. A compound according to Claim 3, named 4-{3-ethyl-5-[(1-ethylpropyl)amino]-6-methoxypyrazin-2-yl}-3-methoxyphenol.

107. A compound according to Claim 3, named 6-ethyl-N-(1-ethylpropyl)-5-(4-isopropoxy-2-methoxyphenyl)-3-methoxypyrazin-2-amine.

108. A compound according to Claim 3, named 4-[3,6-diethyl-5-(1-isopropyl-2-methylpropoxy)pyrazin-2-yl]-3-ethylphenol.

109. A compound according to Claim 3, named 3-ethyl-N-(1-ethylpropyl)-5-[4-fluoro-2-(trifluoromethyl)phenyl]-6-methoxypyrazin-2-amine.

110. A compound according to Claim 3, named 2,5-diethyl-3-(1-isopropyl-2-methylpropoxy)-6-mesitylpyrazine.

111. A compound according to Claim 3, named 2,5-diethyl-3-(1-ethylpropoxy)-6-(4-methoxy-2-methylphenyl)pyrazine.

112. A compound according to Claim 3, named 2,5-diethyl-3-[4-methoxy-2-(trifluoromethoxy)phenyl]-6-(1-propylbutoxy)pyrazine.

113. A compound according to Claim 3, named 6-chloro-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine.

114. A compound according to Claim 3, named 2-(4-butoxy-2-chlorophenyl)-3,6-diethyl-5-(1-ethylbutoxy)pyrazine.

115. A compound according to Claim 3, named 3-bromo-N-(1-ethylpropyl)-6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-amine.

116. A compound according to Claim 3, named 2-(4-{{tert-butyl(dimethyl)silyl}oxy}-2-ethoxyphenyl)-3,6-diethyl-5-(1-ethylpropoxy)pyrazine.

117. A compound according to Claim 3, named 6-methoxy-5-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazin-2-ylformamide.

118. A compound according to Claim 3, named 2,5-diethyl-3-(2-ethylbutyl)-6-[2-methoxy-4-(trifluoromethoxy)phenyl]pyrazine.

119. A compound according to Claim 3, named 2,5-diethyl-3-(1-ethylpropoxy)-6-(2-isopropoxy-4-propoxyphenyl)pyrazine.

120. A compound according to Claim 3, named 2-(sec-butylthio)-5-(2,4-dimethoxyphenyl)-3,6-diethylpyrazine.

121. A compound according to Claim 3, named 5-(4-chloro-2,6-dimethoxyphenyl)-6-methoxypyrazin-2-amine.

122. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine.

123. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(2-methyl-4-fluorophenyl)pyrazin-2-amine.

124. A compound according to Claim 3, named 3,6-diethyl-N,N-(2-methoxyethyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine.

125. A compound according to Claim 3, named 3,6-diethyl-N-propyl-N-(cyclopropylmethyl)-5-(2,4-dichlorophenyl)pyrazin-2-amine.

126. A compound according to Claim 3, named 3,6-diethyl-2-(1-ethylpropoxy)-5-(2-methoxy-4,6-dimethylphenyl)pyrazine.

127. A compound according to Claim 3, named 3,6-dimethyl-2-(1-ethylpropoxy)-5-(2,4-dichlorophenyl)pyrazine.

128. A compound according to Claim 3, named 3,6-diethyl-2-(1-ethylpropoxy)-5-(2-hydroxy-4,6-dimethylphenyl)pyrazine.

129. A compound according to Claim 3, named 3,6-diethyl-2-(1-ethylpropoxy)-5-(2,4-dichlorophenyl)pyrazine.

130. A compound according to Claim 3, named 3,6-dimethyl-N-(1-methoxy-2-butyl)-5-(2,4-dimethoxyphenyl) pyrazin-2-amine.

131. A compound according to Claim 3, named 2-(2,4-dichlorophenyl)-5-[2-(methoxymethyl)pyrrolidin-1-yl]-3,6-dimethylpyrazine.

132. A compound according to Claim 3, named 2-(2,4-dimethoxyphenyl)-5-[2-(methoxymethyl) pyrrolidin-1-yl]-3,6-dimethylpyrazine.

133. A compound according to Claim 3, named 3,6-dimethyl-N-(1-ethylpropyl)-5-(2-methyl-4-chlorophenyl)pyrazin-2-amine.

134. A compound according to Claim 3, named 3,6-dimethyl-N-(1-ethylpropyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl)pyrazin-2-amine.

135. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(2-trifluoromethoxy-4-methoxyphenyl)pyrazin-2-amine.

136. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxy-4-trifluoromethylphenyl)pyrazin-2-amine.

137. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(2-trifluoromethyl-4-methoxyphenyl)pyrazin-2-amine.

138. A compound according to Claim 3, named 3,6-diethyl-N-(1-ethylpropyl)-5-(2-methoxy-4-methylphenyl)pyrazin-2-amine.

139. A compound according to Claim 3, named 3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2-chloro-4-dimethylaminophenyl)pyrazin-2-amine.

140. A compound according to Claim 3, named 3,6-diethyl-N-(1-methylpropyl)-5-(2-chloro-4-dimethylaminophenyl)pyrazin-2-amine.

141. A compound according to Claim 3, named 3,6-diethyl-N-(2-methylpropyl)-5-(2-chloro-4-dimethylaminophenyl)pyrazin-2-amine.

142. A compound according to Claim 3, named 3,6-diethyl-N-(2-phenylethyl)-5-(2-chloro-4-dimethylphenyl)pyrazin-2-amine.

143. A compound according to Claim 3, named 3,6-diethyl-N-(1-propylbutyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl) pyrazin-2-amine.

144. A compound according to Claim 3, named 3,6-diethyl-N-(1-methoxy-2-butyl)-5-(2-trifluoromethyl-4-dimethylaminophenyl) pyrazin-2-amine.

145. A compound according to Claim 3, named 2-(2,6-dimethoxy-4-chlorophenyl)-6-ethyl-5-(1-ethylpropoxy)-3-methoxypyrazine.

146. A compound named 2-(2-methoxy-5-trifluoromethoxyphenyl)-3-ethyl-6-methylamino-5-(1-ethylpropoxy)-pyrazine.

147. A method for treating a CNS disorder or disease, comprising
5 administering to a patient in need of such treatment between 0.1 to 140 mg per kg body weight per day of a compound according to Claim 3.

148. The method of Claim 147 wherein the CNS disorder or disease is an anxiety disorder, stress-related disorder, or eating disorder.

10

149. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of a compound according to Claim 3.

150. A compound according to Claim 3 wherein, in a standard in vitro CRF
15 receptor binding assay the compound exhibits an IC_{50} value less than or equal to 1 micromolar.

151. A method for modulating CRF receptor function, comprising
20 administering to a patient in need of such modulation an effective amount of a compound of according to Claim 3.

152. A method for treating a CNS disorder or disease, comprising
administering to a patient in need of such treatment an effective amount of a compound according to Claim 3.

25

153. The method of Claim 152 wherein the CNS disorder or disease is an anxiety disorder, stress-related disorder, or eating disorder.

154. A compound according to Claim 3, wherein in a standard in vitro sodium channel functional assay the compound does not show any statistically significant activity at the $p < 0.05$ level of significance.

5 155. A method of Claim 152 wherein the CNS disease or disorder is depression or a bipolar disorder.

156. A method of Claim 152 wherein the CNS disease or disorder is anorexia nervosa, bulimia nervosa, or obesity.

10

157. A method according to Claim 152 wherein in a standard in vitro CRF receptor binding assay the compound exhibits an IC_{50} value less than or equal to 1 micromolar.

15 158. A method according to Claim 152 wherein in a standard in vitro CRF receptor binding assay the compound exhibits an IC_{50} value less than or equal to 100 nanomolar.

20 159. A method according to Claim 152 wherein in a standard in vitro CRF receptor binding assay the compound exhibits an IC_{50} value less than or equal to 10 nanomolar.

160. A method according to Claim 152 wherein the patient is a human.

25 161. A method for localizing CRF receptors is tissue section samples comprising:

 contacting with a sample of tissue a detectably-labelled compound of Claim 3 under conditions that permit binding of the compound to CRF receptors within the sample of tissue;

30 washing the tissue sample to remove unbound compound; and

detecting remaining bound compound.

162. A packaged pharmaceutical composition comprising a pharmaceutical composition of claim 149 in a container and instructions for using the composition to
5 treat a patient suffering from an anxiety disorder, a stress-related disorder, or an eating disorder.

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number
WO 01/60806 A3

(51) International Patent Classification⁷: **C07D 241/18**,
241/20, 241/12, 409/04, 401/04, 403/04, 401/12, 405/12,
A61K 31/4965, A61P 25/00, C07D 241/52

(74) Agent: **CORLESS, Peter, F.**: Edwards & Angell LLP,
Dike, Bronstein, Roberts & Cushman, Intellectual Property
Group, 130 Water Street, Boston, MA 02109 (US).

(21) International Application Number: **PCT/US01/05264**

(22) International Filing Date: 16 February 2001 (16.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/182,934 16 February 2000 (16.02.2000) US
60/206,455 22 May 2000 (22.05.2000) US

(71) Applicant: **NEUROGEN CORPORATION** [US/US]: 35
Northeast Industrial Road, Branford, CT 06405 (US).

(71) Applicants and

(72) Inventors: **YOON, Taeyoung** [KR/US]: 6 Finch Lane,
Guilford, CT 06437 (US). **GE, Ping** [CN/US]: 718R
Higganum Road, Durham, CT 06422 (US). **HORVATH,**
Raymond, F. [US/US]: 973 Little Meadow, Guilford, CT
06437 (US). **DE LOMBAERT, Stephane** [BE/US]: 37
Concord Drive, Madison, CT 06443 (US). **HODGETTS,**
Kevin, J. [GB/US]: 224 Reservoir Road, Killingworth, CT
06419 (US). **DOLLER, Dario** [US/US]: 20 Killen Road,
Wallingford, CT 06492 (US). **ZHANG, Cunyu** [CN/US]:
109 Montoya Drive, Branford, CT 06405 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

with international search report

(88) Date of publication of the international search report:
7 February 2002

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **SUBSTITUTED ARYLPYRAZINES**

(57) Abstract: Arylpyrazine compounds are provided, including arylpyrazines that can bind with high affinity and high selectivity to CRF₁ receptors, including human CRF₁ receptors. The invention thus includes methods for treatment of disorders and diseases associated with CRF₁ receptors, including CNS-related disorders and diseases, particularly affective disorders and diseases, and acute and chronic neurological disorders and diseases.

WO 01/60806 A3

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 01/05264

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D241/18 C07D241/20 C07D241/12 C07D409/04 C07D401/04 C07D403/04 C07D401/12 C07D405/12 A61K31/4965 A61P25/00 C07D241/52		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 38174 A (GLAXO GROUP LTD.) 3 September 1998 (1998-09-03) cited in the application the whole document	1-162
Y	WO 95 10506 A (THE DU PONT MERCK PHARMACEUTICAL CO.) 20 April 1995 (1995-04-20) the whole document	1-162
A	WO 98 11075 A (THE DU PONT MERCK PHARMACEUTICAL CO.) 19 March 1998 (1998-03-19) the whole document	1-162
P, A	WO 00 12488 A (GLAXO GROUP LTD.) 9 March 2000 (2000-03-09) the whole document	1-162
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*&* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search <div style="text-align: center; font-weight: bold; font-size: 1.2em;">17 September 2001</div>		Date of mailing of the international search report <div style="text-align: center; font-weight: bold; font-size: 1.2em;">21/09/2001</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer <div style="text-align: center; font-weight: bold; font-size: 1.2em;">Beslier, L</div>

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/05264

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9838174	A	03-09-1998	AU 732915 B2	03-05-2001
			AU 6823798 A	18-09-1998
			BG 103723 A	31-05-2001
			BR 9807814 A	22-02-2000
			CN 1253551 T	17-05-2000
			EE 9900376 A	17-04-2000
			WO 9838174 A1	03-09-1998
			EP 0966448 A1	29-12-1999
			HR 980107 A1	31-12-1998
			HU 0001802 A2	28-05-2001
			JP 2000511203 T	29-08-2000
			NO 994213 A	29-10-1999
			PL 335441 A1	25-04-2000
			SK 117399 A3	12-06-2000
			TR 9902082 T2	21-04-2000
			US 6255307 B1	03-07-2001
WO 9510506	A	20-04-1995	AU 692484 B2	11-06-1998
			AU 8012294 A	04-05-1995
			BR 9407799 A	06-05-1997
			CA 2174080 A1	20-04-1995
			CN 1142817 A	12-02-1997
			CZ 9601014 A3	13-11-1996
			EP 0723533 A1	31-07-1996
			FI 961599 A	07-06-1996
			HR 940664 A1	31-12-1996
			HU 74464 A2	30-12-1996
			JP 9504520 T	06-05-1997
			NO 961425 A	12-06-1996
			NZ 274978 A	27-04-1998
			PL 313973 A1	05-08-1996
			SK 47096 A3	01-10-1996
			WO 9510506 A1	20-04-1995
			US 6107301 A	22-08-2000
			ZA 9407921 A	11-04-1996
WO 9811075	A	19-03-1998	AU 4270297 A	02-04-1998
			EP 0927171 A1	07-07-1999
			JP 2001502300 T	20-02-2001
			WO 9811075 A1	19-03-1998
			US 6218391 B1	17-04-2001
			US 6159980 A	12-12-2000
WO 0012488	A	09-03-2000	AU 5624999 A	21-03-2000
			BR 9913183 A	15-05-2001
			WO 0012488 A1	09-03-2000
			EP 1107960 A1	20-06-2001

3
-
1

THIS PAGE BLANK (USPTO)
